

## www.ancienttl.org · ISSN: 2693-0935

Ancient TL, 2006. *Thesis Abstracts*. Ancient TL 24(1): 23. https://doi.org/10.26034/la.atl.2006.392

This article is published under a *Creative Commons Attribution 4.0 International* (CC BY): https://creativecommons.org/licenses/by/4.0



© Ancient TL, 2006

## **Thesis Abstract**

Author:	Sara Ante
Thesis Title:	Luminescence dating tests of
	detrital grains from sediment
	traps, Andvord Bay and
	Brialmont Cove, Antarctic
	Peninsula
Grade:	MSc
Date:	December 2005
Supervisor:	Glenn Berger
Address:	University of Nevada-Reno,
	Department of Geological
	Sciences and Engineering,
	United States of America, and
	Desert Research Institute, Reno
	(Berger)

Difficulties (e.g., large, diverse, erratic reservoir corrections) with radiocarbon dating have prompted photonic dating tests on Antarctic glacial marine sediments. This technique (clock zeroing is based on the last exposure to daylight of quartz and feldspar grains) was applied to modern-age sediments collected in two well-characterized fjords on the western side of the Antarctic Peninsula. Our deployed sediment traps collected approximately 14 months of deposition between January 2002 and March 2003. We conducted a series of multi-aliquot infrared-stimulated luminescence (IRSL) tests on polymineral silt-sized grains to evaluate the extent of the clock-zeroing process in this setting, to provide a foundation for applications elsewhere in the region. Comparisons were made between samples on the basis of seasonal differences and trap depth. Several double single-aliquot-regenerative (SAR) dose experiments were also performed on one of the sediment trap samples. Sediments representing summer and winter deposition were also characterized by microprobe analyses.

Had the sediments been exposed to sufficient sunlight at deposition, sediment grains collected by these traps would exhibit very low equivalent-dose ( $D_E$ ) values. However, single and multi-aliquot experiments resulted in high (and variable)  $D_E$  values for both Andvord Bay and Brialmont Cove, which yield erroneously old ages, rather than zero. IRSL experiments have resulted in apparent ages for Andvord Bay ranging between 19.8 ± 1.4 ka and 56.8 ± 5.4 ka ( $D_E = 48 - 123$  Gy), while Brialmont Cove  $D_{\rm E}$  values ranged between 11.1  $\pm$  0.7 ka and 31.4  $\pm$ 3.4 ka ( $D_E = 21 - 58$  Gy), depending on season and depth of deposition. This indicates that these glacial marine sediments were unevenly and poorly bleached Variations in luminescence traits at deposition. between seasons and with trap depth may, however, be helpful for studying sedimentation processes in these fjords. Double-SAR experiments also resulted in anomalously high D<sub>E</sub> values. SAR data for one sediment trap sample did exhibit characteristics predicted by Bailey et al.'s (2003)  $D_E(t)$  model for poorly bleached sediments. Microprobe analyses did not reveal any significant differences in mineralogy between seasons and depth that would affect apparent age results.