

# www.ancienttl.org · ISSN: 2693-0935

Issue 6(3) - December 1988 https://doi.org/10.26034/la.atl.v6.i3

This issue is published under a Creative Commons Attribution 4.0 International (CC BY): https://creativecommons.org/licenses/by/4.0



© Ancient TL, 1988

# Irradiation of loess samples at elevated temperatures

### A.G. Wintle\* and S.C. Packman<sup>†</sup>

\*Geography Department, Royal Holloway and Bedford New College, Egham, Surrey, TW20 0EX, UK.

<sup>†</sup>Godwin Laboratory, Free School Lane, Cambridge CB2 3RS, UK.

#### Introduction

In some recent TL studies of loess from western Europe, it has not been possible to achieve ages greater than about 100 ka (Debenham, 1985) or 150 ka (Wintle, Shackleton and Lautridou, 1985; Wintle, 1985). Systematic underestimation of the TL ages occurs when the TL from mixed fine grain minerals is observed in the ultraviolet part of the spectrum, as observed using a Schott UG-11 filter, and when the regeneration method is used for ED determination for samples older than about 50 ka. Typical European loess samples of this age will have an ED of around 200 Gy and their natural TL signal will occur in a part of the glow curve which is clearly non-linear. It should be pointed out that no such underestimation has been reported by Zöller, Stremme and Wagner, 1988), who employ the regeneration method on fine grains but use a Corning 5-58 filter and a strong thermal wash before TL measurement.

Several causes have been hypothesized, but none proven.

1) The apparent saturation of the TL age obtained for progressively older samples led Debenham (1985) to suggest a time-dependent loss of luminescence centres. This would effect all methods of ED determination, not just the regenerative method.

2) A sensitivity change after bleaching. No sensitivity changes are apparent for young samples, and it is difficult to determine such changes for older samples which show non-linear growth curves. However, it has been hypothesized that the sensitivity change is dose-dependent (Wintle, 1985; Rendell and Townsend, 1988).

#### The hypothesis

There is also another difference between irradiation in nature which produces the natural TL signal and irradiation of the laboratory-

bleached sample discs which results in the matching regenerated signal - the traps which are responsible for the TL signal up to 200 °C are kept empty at ambient temperature in the natural environment. Let us hypothesize that the trapping cross section for these traps is higher than for the traps between 280 °C and 380 °C, where the ED determinations are made. Then during natural irradiation free charge carriers will go preferentially to the shallower traps and the deeper traps will fill more slowly than during the laboratory irradiation after bleaching when the shallower traps fill and stay filled. This would result in an apparent increase in TL sensitivity after bleaching and thus an underestimated ED.

To see whether this is occurring irradiations need to be carried out at elevated temperatures. In the literature there have been several instances of sensitivity changes being reported as a result of irradiations at different temperatures; for example for LiF (TLD-100) (Chandra et al, 1982). Aitken et al (1974) found a 40% decrease in the TL signal for the 325 °C peak in quartz when irradiated at 200 °C compared with irradiation at 20 °C. Durrani et al (1977) found a large decrease in the intensity of the TL signal from several quartzes as they decreased the irradiation temperature from room temperature to 113 K.

#### Experiment

A tube oven was placed in the beam of the  ${}^{60}$ Co gamma source in the Biochemistry Department in Oxford. Five discs (fine grains deposited onto 1 cm aluminium discs in the usual way) were placed in identical positions in an aluminium container in the oven for each temperature. The container was allowed to reach equilibrium temperature before the irradiation was performed. Temperatures of 18, 60, 100 and 123 °C were used. An irradiation time of one hour, giving about 130 Gy, was used for each experiment. The discs were unfortunately

not able to be measured until 8 weeks after the irradiation. The results are shown in figure 1 and show no difference in the TL signal above 270 °C. As well as irradiating bleached discs, a comparison was made between irradiation at 18 and 100 °C using the same dose but adding it on top of the natural dose. Again no difference was observed.

#### Conclusion

The similarity of the glow curves above 270 °C does not confirm the hypothesis outlined in the previous section.

#### References

- Aitken, M.J., Huxtable, J., Wintle, A.G., and Bowman, S.G.E. (1974) Age determination by TL: review of progress at Oxford. Fourth Int. Conf. on Lumin. Dosim., Krakow, Poland, 1005-1019.
- Chandra, B., Lakshmanan, A.R., and Bhatt, R.C. (1982) Dependence of thermoluminescence sensitivity on temperature of irradiation in LiF (TLD-100) phosphor. J. Appl. Phys. D, 15, 1803-1816.
- Debenham, N.C. (1985) Use of UV emissions in TL dating of sediments. *Nucl. Tracks and Radn. Measts.*, **10**, 717-724.

- Durrani, S.A., Groom, P.J., Khazal, K.A.R., and McKeever, S.W.S. (1977) The dependence of the thermoluminescence sensitivity upon the temperature of irradiation of quartz. J. Phys. D, 10, 1351-1361.
- Rendell, H.M., and Townsend, P.D. (1988) Thermoluminescence dating of a 10m loess profile in Pakistan. *Quat. Sci. Revs.* 7(3&4), 251 - 255.
- Wintle, A.G. (1985) Sensitization of TL signal by exposure to light. Ancient TL, 3, 16-21.
- Wintle, A.G., Shackleton, N.J., and Lautidou, J.P.(1984) Thermoluminescence dating of periods of loess deposition and soil formation in Normandy. *Nature*, 310, 491-493.

ŝ

Zöller, L., Stremme, H., and Wagner, G.A. (1988) Thermolumineszenz-Datierung an Loess-Palaoboden-Sequenzen von Nieder-, Mittel- und Oberrhein/-Bundersrepublik Deutschland. *Chemical Geology* (Isotope Geoscience Section), 73, 39-62. (In German).

PR Reviewed by S.G.E. Bowman



Figure 1. Glow curves of fine grains for a sample of Hungarian loess (QTL85DB) which had been bleached and then given a gamma dose of 130 Gy at temperatures of a) 18 °C, b) 60 °C, c) 100 °C, and d) 123 °C. Measurement was with a Schott UG-11 filter at a heating rate of 5 %.

## Bibliography

- Bouchez, R., Ma, J-L, Amosse, J., Braundet, J-F, and Piboule, M. (1987) Resonance paramagnetique clectronique. In *Geologie de la prehistoire; methodes, techniques, applications* (Ed. J-C Miskovsky), 1111-1123. (in French).
- Cini-Castagnoli, G. (1988) Stability of the cyclic behaviour of the sun in the past millenia. In Solarterrestrial Relationships and the earth environment in the last millenia XCV Corso, 317-328.
- Cini-Castagnoli, G., and Bonino, G. (1988) Solar imprint in sea sediments: the thermoluminescence profile as a new proxy record. In Secular, solar and geomagnetic variations in the last 10 000 years (Eds F.R. Stephenson and A.W. Wolfendale), 341-347.
- Cini-Castagnoli, G., Bonino, G., and Provenzale, A. (1988) On the thermoluminescence profile of an Ionian Sea sediment: evidence of 137, 118, 12.1, and 10.8 year cycles in the last two millennia. *Il Nuovo Cimento*, 11C, 1-12.
- Fedorowicz, S. (1988) Thermoluminescence dating of sands. Przegl. Geol., 36, 148-150. (In Polish).
- Forman, S.L. (1988) The solar resetting of thermoluminescence of sediments in a glacier-dominated fiord environment in Spitsbergen: geochronologic implications. Artic and Alpine Research, 20, 243-253.
- Hornyak, W.F., and Franklin, A.D. (1988) A second generation 3-D spectrophotometer for TL analysis. J. Lumin., 42, 89-96.
- Juvigné, E.H., and Wintle, A.G. (1988) A new chronostratigraphy of the Late Weichselian loess units in Middle Europe based on thermoluminescence dating. *Eiszeitalter und Gegenwart*, 38, 94-105.
- Kirsh, Y., Townsend, P.D., and Shoval, S. (1987) Local transitions and charge transport in thermoluminescence of calcite. Nucl. Tracks and Radn. Measts., 13, 115-119.
- Mazumdar, P.S., Singh, S.J., and Gartia, R.K. (1988) The determination of activation energy from the shape of a thermoluminescence peak. J. Phys. D; Appl. Phys., 21, 815-819.
- Molodkov, A. and Raukas, A. (1988) The age of Pleistocene marine deposits of Arkhangelsk based on EPR dating of subfossil mollusc shells. *Izvestiya Akademii Nauk Estonskoy SSR Geologiya*, 37, 36-42. (In Estonian).
- Moscovitch, M., and Horowitz, Y.S. (1988) A microdosimetric track interaction model applied to alphaparticle-induced supralinearity and linearity in thermoluminescent LiF:Mg,Ti. J. Phys. D; Appl. Phys., 21, 804-814.
- Singh, T., Chandra, S., Mazumdar, P.S., and Gartia, R.K. (1988) On the determination of activation energy in thermoluminescence by the initial rise method. J. Phys. D; Appl. Phys., 21, 1312-1314.
- Valladas, H. (1987) Thermoluminescence. In Geologie de la prehistoire; methodes, techniques, applications (Ed J-C Miskowsky), 1061-1071. (In French).
- Zöller, L., Stremme, H., and Wagner, G.A. (1988) Thermolumineszenz-Datierung an Loess-Palaoboden-Sequenzen von Nieder-, Mittel- und Oberrhein. *Chemical Geology* (Isotope Geoscience Section), 73, 39-62. (in German).

#### TL dates quoted in:

Jensen, K.A., and Knudsen, K.L. (1988) Quaternary foraminiferal stratigraphy in boring 81/29 from the central North Sea. *Boreas*, 17, 273-287.

Shafer, D.S. (1988) Late Quaternary landscape evolution at Flat Laurel Gap, Blue Ridge Mountains, North Carolina. Quat. Res., 30, 7-11.

*New in paperback* - Thermoluminescence of Solids by S.W.S. McKeever. Cambridge University Press. Price £17.50 or \$34.50.

Compiled by Ann Wintle