
Ancient TL

www.ancienttl.org · ISSN: 2693-0935

Issue 8(1) - March 1990

<https://doi.org/10.26034/la.atl.v8.i1>

This issue is published under a Creative Commons Attribution 4.0 International (CC BY):

<https://creativecommons.org/licenses/by/4.0>



© Ancient TL, 1990

Regression analysis of exponential palaeodose growth curves

V. Poljakov and G. Hütt

Institute of Geology, Estonian Academy of Sciences, Estonia pst.7, Tallinn 200101, ESSR, USSR.

When using palaeodosimetric dating methods, laboratory reconstruction of the palaeodose poses a key problem. As a rule, it is solved by the *additive dose* method, with extrapolation of the resulting experimental curve towards intersection with the x-axis. Given an experimental set of data (1)

$$\{D_A^{(i)}, I_i \equiv I(D_A^{(i)}); i = 1, \dots, n\} \quad (1)$$

where D_A is the laboratory additive dose, and $I(D_A)$ the intensity of the TL peak or ESR spectrum signal, reconstruction of the accumulated dose D_N is realised through a statistical regression analysis. This assumes optimum choice of the regression functional dependence.

Supposing that fading involves a first order thermally activated process, the solving of the kinetic equations leads to the following functional shape for the intensity-dose relation (Hütt and Smirnov, 1982):

$$I(D_A) = I_0 [1 - e^{-\beta(D_N + D_A)}] \quad (2)$$

where I_0 and β are the parameters characterising the palaeo-dosimeter. Thus, (1) and (2) represent a non-linear three-parameter regression model I_0, β, D_N are the parameters to be estimated.

It would be more convenient to perform the regression analysis rewriting the equation (2) in the following way:

$$y(x) = a + be^{cx}$$

where $x \equiv D_A$ and $y(x) \equiv I(D_A)$.

From the condition $y(-D_N) = 0$ we obtain the value of the accumulated dose:

$$D_N = \frac{1}{c} \ln \left(-\frac{b}{a} \right)$$

The final model may be expressed as follows

$$y_i = f(\vec{\theta}; x_i) + \varepsilon_i, \quad i = 1, \dots, n$$

where y_i is the dependent variable, $\vec{\theta} = \{a, b, c\}$ is the estimated parameter vector, x_i is the independent variable, and $\vec{\varepsilon}$ is the random deviation vector. It is also assumed that x_i is error-free and that the random deviations are of normal distribution and uncorrelated,

i.e. $\text{cov}(\vec{\varepsilon}) \sim N(\vec{0}, \sigma^2 \vec{I})$.

The parameters a, b, c are estimated using the least squares method, when optimal choice of these parameters $\hat{\vec{\theta}} = \{\hat{a}, \hat{b}, \hat{c}\}$ is determined by minimization of

$$S(a, b, c; x_i, y_i) = \sum_{i=1}^n (y_i - a - be^{cx_i})^2$$

The linearised Newton-Gauss method of minimization of $S(\vec{\theta}; x_i, y_i)$ is known from the literature (Berger et al, 1987). However, the linearisation procedure has some disadvantages. In some cases it leads to a slow convergence of the iterative processes and even to divergence.

The present paper proposes a straight minimisation method of $S(\vec{\theta}; x_i, y_i)$ by solving of the corresponding system of normal equations:

$$\left. \frac{\partial S}{\partial a} \right|_{\vec{\theta} = \hat{\vec{\theta}}} = - \sum_{i=1}^n 2 (y_i - \hat{a} - \hat{b}e^{\hat{c}x_i}) = 0$$

$$\left. \frac{\partial S}{\partial b} \right|_{\vec{\theta} = \hat{\vec{\theta}}} = - \sum_{i=1}^n 2 (y_i - \hat{a} - \hat{b}e^{\hat{c}x_i}) e^{\hat{c}x_i} = 0$$

$$\left. \frac{\partial S}{\partial b} \right|_{\vec{\theta} = \hat{\vec{\theta}}} = - \sum_{i=1}^n 2 (y_i - \hat{a} - \hat{b}e^{\hat{c}x_i}) \hat{b}x_i e^{\hat{c}x_i} = 0 \quad \dots (3)$$

From the first two equations of the system (3) we have:


$$\hat{b} = \frac{\sum_{i=1}^n y_i e^{\hat{c}x_i} - \frac{1}{n} \sum_{i=1}^n e^{\hat{c}x_i} \sum_{i=1}^n y_i}{\sum_{i=1}^n e^{2\hat{c}x_i} - \frac{1}{n} \left(\sum_{i=1}^n e^{\hat{c}x_i} \right)^2}$$

$$\hat{a} = \frac{1}{n} \sum_{i=1}^n y_i - \frac{b}{n} \sum_{i=1}^n e^{\hat{c}x_i} \quad \dots (4)$$

Inserting (4) into the third equation of the system (3) we obtain the equation for $\hat{c}$:

$$F(\hat{a}, \hat{b}, \hat{c}; x_i, y_i) = 0 \quad (5)$$

Non-linear equation (5) can be solved numerically by means of the Newton-Raphson iteration method:



$$\hat{c}_K = \hat{c}_{K-1} - \frac{F(\hat{c}_{K-1})}{F'(\hat{c}_{K-1})}, \quad K = 1, \dots; \quad (6)$$

where $\hat{c}_0$ is the initial approximation and

$$F(c) \equiv \frac{\partial F}{\partial c} + \frac{\partial F}{\partial a} \frac{\partial a}{\partial c} + \frac{\partial F}{\partial b} \frac{\partial b}{\partial c}$$

Calculating the derivatives included in $F(\hat{c}_{K-1})$ by means of the system of equations (4) we entirely determine the calculation scheme of the iterative procedure. The termination of the iterative process (6) is based on the natural proximity condition of consequent estimates for $\hat{c}$:

$$|(\hat{c}_{K*}) - (\hat{c}_{K-1})| \leq \varepsilon$$

The proposed calculation scheme is realised in the form of a BASIC programme. The working results of the programme are presented in the figure, where the regression dependence is shown by the solid line, and the experimental data by dots.

A linearised procedure for the approximate estimation of error in the accumulated dose can be carried out in the following way. Denoting with $V(y_i)$ the variance of y_i :

$$V(y_i) = \frac{1}{m_i - 1} \sum_{j=1}^{m_i} (y_{ij} - \bar{y}_i)^2 \quad i = 1, \dots, n;$$

$$\text{where } \bar{y}_i = \frac{1}{m_i} \sum_{j=1}^{m_i} (y_{ij} - y_i)^2 \quad i = 1, \dots, n;$$

and m_i is the number of repeated measurements of value y_i . Writing the accumulated dose in a Taylor series in powers of $\delta y_i \equiv y_i - \bar{y}_i$, and confining ourselves to linear terms:

$$\delta D_N = \sum_{i=1}^n \left(\frac{\partial f}{\partial a} \frac{\partial a}{\partial y_i} + \frac{\partial f}{\partial b} \frac{\partial b}{\partial y_i} + \frac{\partial f}{\partial c} \frac{\partial c}{\partial y_i} \right) \delta y_i, \quad (7)$$

where

$$f = D_N = \frac{1}{c} \ln \left(\frac{b}{a} \right),$$

$$a = a(x_i, y_i), \quad b = b(x_i, y_i), \quad c = c(x_i, y_i);$$

$$\delta D_N = f(a, b, c) - f(\hat{a}, \hat{b}, \hat{c})$$

and the derivatives of function f are determined at the point of

$$\hat{\theta} = \{ \hat{a}(x_i, \bar{y}_i), \hat{b}(x_i, \bar{y}_i), \hat{c}(x_i, \bar{y}_i) \}.$$

A linearized procedure for the approximation estimate of error in the accumulated dose can be carried out by using the law of propagation of errors (Mandel, 1964):

$$V(D_N) = \sum_{i=1}^n \left(\frac{\partial D_N}{\partial a} \frac{\partial a}{\partial y_i} + \frac{\partial D_N}{\partial b} \frac{\partial b}{\partial y_i} + \frac{\partial D_N}{\partial c} \frac{\partial c}{\partial y_i} \right)^2 V(y_i)$$

where $V(D_N)$ is the variance of D_N and $V(y_i)$ is the variance in the value of y_i evaluated using replicate measurements at the same dose x_i .

References

- Berger, G.W., Lockhart, R.A., and Kuo, I., (1987), Regression and error analysis applied to the dose-response curves in thermoluminescence dating. *Nucl. Tracks and Radn. Meas.*, **13**, 177-184.
- Hutt, G., and Smirnov, A., (1982) Detailed thermoluminescence dating studies of samples from geological reference profiles in Central Russia. *PACT J.*, **6**, 505-513.
- Mandel, J. (1964) *The statistical analysis of experimental data*. Interscience, p73.

PR. Reviewed by A.D. Franklin

Bibliography

The bibliography is devoted to those papers included in the *Proceedings of the 2nd International Symposium on ESR Dosimetry and Applications* which have been published in (1989) *Applied Radiation and Isotopes*, **40** (10-12). (Eds D.F. Regulla, A. Scharmann, and W.L. McLaughlin)

Historical Perspective

Scharmann, A. ESR - a scientific tool for applications, 829.

Methods and Instruments

Franconi, C., Holowacz, J., Ettinger, K. V., Staderini, E.M., Bonori, M., and Laitano, R.F. A precise S-band ESR reader for free-radical routine dosimetry, 835.

Hara, Hidemoto, and Ikeya, Motoji. Frequency-sweep ESR spectrometer for dosimetry and dating, 841.

Ikeya, Motoji, and Furusawa, Masahiro. A portable spectrometer for ESR microscopy, dosimetry and dating, 845.

Kojima, T, and Tanaka, R. Polymer-alanine dosimeter and compact reader, 851.

Correlation with other Solid State Effects

Halliburton, L. E. ESR and optical characterization of point defects in quartz, 859.

Ettinger, K. V. Free radical dosimetry techniques and their suitability for precise and accurate measurements of radiation, 865.

Azorin, J., Gutiérrez, A., Munoz, E., and Gleason, R. Correlation of ESR with lyoluminescence dosimetry using some sugars, 871.

Kundu, H., Sur, T., Mitra, B., and Roy, S.C. Study of decay of trapping centres of free radicals in high-temperature icy states of saccharides by lyoluminescence and ESR methods, 875.

Oommen, I.K., Nambi, K.S.V., Sengupta, S., Gundo Rao, T.K., and Ravikumar, M. Lactose and "Tris" lyoluminescence dosimetry systems and ESR correlation studies, 879.

Dosimetry

Van Laere, K., Buysse, J., and Berkvens, P. Alanine in high-dose dosimetry: spectrophotometric and electro-chemical readout procedures compared with ESR, 885.

Kotzias, D., and Beyerle-Pfnür, R. Characterization of humic substances by ESR spectroscopy, 897.

Simmons, J.A. Thermal effects in irradiated amino acids, 901.

Göksu, H.Y., Wieser, A., Waibel, A., Vogenauer, A., and Regulla, D.F. Comparing measurements of free radicals, optical density and thermoluminescence in solids for high-level dosimetry, 905.

Wieser, A., and Regulla, D.F. ESR dosimetry in the "gigarad" range, 911.

Dosimetry - Response to Neutrons and Heavy Ions

Hüttermann, J., and Schaefer, A. Heavy-iron induced free radical formation in solid DNA-constituents: quantitative and structural aspects, 915.

Waligórski, M.P.R., Daniały, G., Sun Loh, K., and Katz, R. The response of the alanine detector after charged-particle and neutron irradiations, 923.

Hansen, J.W., and Olsen, K.J. Predicting decay in free-radical concentration in L- α -alanine following high-LET radiation exposures, 935.

Schraube, H., Weitzenegger, E., Wieser, A., and Regulla, D. F. Fast neutron response of alanine probes, 941.

Reference Dosimetry

McLaughlin, W.L. Reference dosimetry and measurement quality assurance, 945.

Nam, J.W., and Regulla, D.F. The significance of the International Dose Assurance Service for radiation processing, 953.

Wieser, A., Siegele, R., and Regulla, D.F. Influence of the irradiation on the free-radical response of alanine, 957.

Sollier, T.J.L., Mosse, D.C., Chartier, M.M.T., and Joli, J.E. The LMRI ESR/alanine dosimetry system: description and performance, 961.

Miller, A., Kovacs, A., Wieser, A., and Regulla, D.F. Measurements with alanine and film dosimeters for industrial 10 MeV electron reference dosimetry, 967.

Pańta, P.P., Strzelczak-Burlinska, G., and Tomasinski, Z. ESR/L-alanine system as a proposed standard dosimeter for electron-beam irradiations, 971.

Coninckx, F., Schonbacher, H., Bartlotta, A., Onori, S., and Rosati, A. Alanine dosimetry as the reference dosimetric system in accelerator radiation environments, 977.

Transfer Dosimetry

- Olsen, K.J., Hansen, J.W., and Waligórski, M.P.R. ESR dosimetry in calibration intercomparisons with high-energy photons and electrons, 985.
- Schneider, Manfred K.H. Realization and dissemination of the unit of absorbed dose to water in the dose range of radiation therapy, 989.
- Chu, S., Wieser, A., Feist, H., and Regulla, D.F. ESR/alanine dosimetry of high-energy electrons in radiotherapy, 993.
- Hjortenberget, P., Hansen, J.W., and Wille, M. Measurement of dose rate from $^{106}\text{Ru}/^{106}\text{Rh}$ ophthalmic applicators by means of alanine-polymer foils, 997.
- Bartolotta, A., Onori, and Pantaloni, M. Intercomparison of the dosimetric system used in industrial irradiation plants in Italy, 1003.
- Dorda, E. M. Dose control in the semi-industrial irradiation plant at Ezeiza Atomic Center, 1009.

Radiation Accident Dosimetry

- Dalgarno, R.G., and McClymont, J.D. Evaluation of ESR as a radiation accident dosimetry technique, 1013.
- Ikeya, Motoji and Ishii, Hiroshi. Atomic bomb and accident dosimetry with ESR: natural rocks and human tooth *in-vivo* spectrometer, 1021.
- Barthe, J., Kamenopoulou, V., Cattoire, B., and Portal, G. Dose evaluation from textile fibres: a post-determination of initial ESR signal, 1029.
- Shimano, Tatsuya, Iwasaki, Midori, Miyazawa, Chyuzo, Miki, Toshikatsu, Kai, Ayako, and Ikeya, Motoji. Human tooth dosimetry for gamma-rays and dental x-rays using ESR, 1035.
- Regulla, D.F., and Deffner, U. Dose estimation by ESR spectroscopy at a fatal radiation accident, 1039.

Dating - General Aspects

- Grün, R. Present status of ESR-dating, 1045.
- Hütt, G., and Jaek, J. Dating accuracy from laboratory reconstruction of palaeodose, 1057.
- Lyons, R.G., and Brennan, B.J. Alpha-particle effectiveness in ESR dating: energy dependence and implications for dose-rate calculations, 1063.
- Renyou, Liang, Zicheng, Peng, Sizhao, Jin, and Peihua, Huand. Estimation of the influence of experimental conditions on ESR dating results, 1071.
- Grün, R., and MacDonald, P.D.M. Non-linear fitting of TL/ESR dose-response curves, 1077.

Dating - Carbonates

- Skinner, A. F. ESR dosimetry and dating in aragonitic molluscs, 1081.
- Molodkov, A. The problem of long-term fading of absorbed palaeodose on ESR-dating of Quaternary mollusc shells, 1087.
- Smith, B.W., Smart, P.L., Fox, P.J., Prescott, J.R., and Symons, M.C.R. An investigation of ESR signals and their related TL emission in speleothem calcite, 1095.
- Barabas, M., Bach, A., Mudelsee, M., and Mangini, A. Influence of the Mg-content on ESR-signals in synthetic calcium carbonate, 1105.
- Katzenberger, O., Debuyst, R., De Canniere, P., Dejeht, F., Aspers, D., and Barabas, M. Temperature experiments on mollusc samples: an approach to ESR signal identification, 1113.
- Peihua, Huang, Renyou, Liang, Sizhao, Jin, Zicheng, Peng, and Rutter, Nathaniel W. Study on accumulated dose in littoral shells of Argentina, 1119.
- Yijian, Chen, Fortin, P., and Li, Qiao. A preliminary study on ESR dating of sediments from Curitiba Basin, Brazil, 1123.
- Zicheng, Peng, Sizhao, Jin, Renyoy, Liang, Peihua, Huang, Yucai, Quan, and Ikeya, M. Study on comparison of ESR dating of coral and shells with $^{230}\text{Th}/^{234}\text{U}$ and ^{14}C methods, 1127.
- Rossi, A.M., and Poupeau, G. Radiation-induced paramagnetic species in natural calcite speleothems, 1133.
- Kai, Ayako, and Ikeya, Motoji. ESR study of fossil shells in sediments at Hamana Lake, 1139.
- Arakawa, Tatsuhiko, and Hori, Nobuyuki. ESR dating of carbonate speleothem rings and late Quaternary climatic changes in the Ryukyu Islands, Japan, 1143.
- Jacobs, C., De Canniere, P., Debuyst, R., Dejeht, F. and Apers, D. ESR study of gamma-ray irradiated synthetic calcium carbonates, 1147.
- Lyons, R.G., Crossley, P.C., Ditchburn, R.G., McCabe, W.J., and Whitehead, N. Radon escape from New Zealand speleothems, 1153.
- Özer, A.M., Wieser, A., Göksu, H.Y., Muller, P., Regulla, D.F., and Erol, O. ESR and TL age determination of caliche nodules, 1159.

Dating - Other Minerals and Enamel

- Yijian, Chen, Arakel, A. V., and Jinfen, Lu. Investigation of sensitive signals due to gamma-ray irradiation of chemical precipitates. A feasibility study for ESR dating of gypsum phosphate and calcrete deposits, 1163.
- Toyoda, Shin, and Ikeya, Motoji. ESR as a paleothermometer of volcanic materials, 1171.
- Imai, N., and Shimokawa, K. ESR dating of the tephra "Crystal Ash" distributed in Shinshu, central Japan, 1177.
- Fukuchi, Tatsuro. Theoretical study on frictional heat by faulting using ESR, 1181.
- Desrosiers, M.F., Simic, M.G., Eichmiller, F.C., Johnston, A.D., and Bowen, R.L. Mechanically-induced generation of radicals in tooth enamel, 1195.
- Soo Chong, Tai, Ohta, Hiroyuki, Nakashima, Yoshiyuki, Iida, Takao, Ieda, Kengo, and Saisho, Hideo. ESR dating of elephant teeth and radiation dose rate estimation in soil, 1199.

Identification of Irradiated Food

- Bögl, K.W. Identification of irradiated foods-methods, development and concepts, 1203.
- Dodd, N.J.J., Lea, J.S., and Swallow, A.J. The ESR detection of irradiated food, 1211.
- Raffi, J., Evans, J.C., Agnel, J.P., Rowlands, C.C., and Lesgards, G. ESR analysis of irradiated frogs' legs and fishes, 1215.
- Ikeya, M., Baffa, Folho, O., and Mascarenhas, S. Quality assessment of coffee beans with ESR and gamma-ray irradiation, 1219.

Technical Note

- Troup, G. J., Philbrow, J.R., Hutton, D.R., Hunter, C.R., and Wilson, G.L. EPR detection of free radicals in (I) coffee and (II) gamma-ray irradiated foodstuffs, 1223.

ESR Imaging and Other Applications

- Eaton, G.R., Eaton, S.S., and Maltempo, M.M. Three approaches to spectral-spatial EPR imaging, 1227.
- Hofer, P., Holczer, K., and Schmalbein, D. Characterization of gamma-ray irradiated powder alanine samples by pulsed EPR, 1233.
- Morita, Yasuyuki, Ohno, Keiichi, Ohashi, Kazutoshi, and Sohma, Junkichi. ESR imaging investigation on depth profiles of radicals in organic solid dosimetry, 1237.
- Miki, T. ESR spatial dosimetry using localized magnetic field modulation, 1243.

Notices

E-mail Address update

Michael Short of the Radioisotope Unit, Hong Kong writes to inform AnTL subscribers that the Laboratory may now be reached via BITNET using the following address:
 HRXRSMA@HKUCC.BITNET

Meetings

6th International Specialist Seminar on TL and ESR Dating. 2-6 July 1990, Clermont Ferrand, France.
 Further details: TL+ESR, Lab. de Physique Corpusculaire, F-63177 Aubiere, Cedex, France.

5th Nordic Conference on the Application of Scientific Methods in Archaeology. Stockholm 20-24 September 1990.

Further details: Stockholms Universitet, Arkeologiska Forskningslaboratoriet, Greens Villa, S-106 91 Stockholm.

Time and Environment - A PACT Seminar, Helsinki, Finland September. 25-28 1990
 Further details: The Dating laboratory, University of Helsinki, Snellmaninkatu, Finland.

NB All of the above meetings are now in the final stages of preparation.

Ancient TL SUPPLEMENT

Date List

March 1990 Issue 3

1. This list includes dates for fired materials of archaeological interest, submitted to *Ancient TL* during 1989 for which sufficient information has been supplied. Readers are referred to earlier issues of the Date List for a fuller description of the structure of entries.
2. Application forms are available from the Editor, who will be pleased to advise on data compilation; laboratories wishing to submit dates for which the current date entry specification is not suitable should write to him. The application forms may be supplied on either paper or magnetic media.
3. A separate summary of all Part 1 entries published so far will be available for general archaeological circulation during 1990.
4. A specification for *Luminescence Sediment Age* entries is now being formulated. Suggestions for the form of the Entry Specification are welcomed and should be sent to the Editor by 1st May 1990.

Laboratory: [name]

Date Entry Specification

Entry: [entry number]

PART I

Site: [Name] Location: [Region, country] Grid Ref.: [National map reference]

Site Description: [Brief description of period and nature of site]

Dates/Ages:

Lab. Ref. Material Archaeological Ref.

[Type]	[Type]	[Lab. abbrev.]					
TL	Context	Date	800	AD ± 50	(Dur87TLfg)	100-1/6	pottery ABC-1a
	Single	Age					
			[Overall error]	[Test year]	[Technique]	[Sample ref.]	[Context reference]
						[Dated material]	

— TL Context Components: [Details of component TL dates/ages used to derive Context Date/Age] —

Archaeological Evidence: [Excavator's brief description of context(s)]

Site Director: [Full name and institutional postal address]

Reports: [Details of excavation and laboratory reports]

PART II

Section A. TL Measurements

1. min.([mineral]) tech.([technique]; [grain size range, gsr] μm)
Data tabulated for each sample:
2. $P = [\text{value}] \pm \text{s.e. Gy}$ 2a. $I/P = [\text{value}]$ 3. Slopes [2nd/1st: $[\text{value}] \pm \text{s.e.}$]
4. [Type of plateau] Plateau $[\pm [\text{value}] \%; [T_1 - T_2]]$
- 4a. Peak [@ $[\text{value}] ^\circ\text{C}$; [heating rate] $^\circ/\text{s}$; [pre-heat details if applicable]]
5. Stability[interval, $T_1 - T_2$]; [period] ; [storage $T ^\circ\text{C}$] ; [result ; $[\text{value}] \pm [\text{value}] \%$]
6. a value = $[\text{value}]$, or b value = $[\text{value}]$

Section B. Dose-rate Measurements

Data tabulated for each sample:

1. Total Effective Dose-rate = $[\text{value}] \pm \text{s.e. mGy/a}$ [$\alpha = [\text{value}] \%$ [method] ;
 $\beta = [\text{value}] \%$ [method] ; $\gamma = [\text{value}] \%$ [method] ; $\cos(\text{mic}) = [\text{value}] \%$ [method]]
2. Radon $[\pm [\text{value}] \%$ [method]]
3. Water [Sample ($[\text{value}] \pm \text{s.e.} \%$); (Burial) Environment ($[\text{value}] \pm \text{s.e.} \%$)]

Section C. Error [[Procedure : eA76 or specify other]]

Section D. TL Age

Data tabulated for each sample:

TL Age $[\pm [\text{random error}] ; \pm [\text{overall accuracy}]]$

Special Remarks: [Details of entries
with * or any other additional information]

KEY TO ABBREVIATIONS

STANDARD METHODS/TECHNIQUES/PROCEDURES

i	Inclusion	pd	Pre-dose	a Plat	Age plateau
fg	Fine-grain	MA	Multiple activation	d Plat	Dose plateau
mmi	Multi-mineral	ADD	Additive dose proc.	s Plat	TL Signal plateau
		Sb	Sensitivity baseline		
α -c	Alpha counting	FPh	Flame Photometry	TLD	TL dosimetry
AAS	Atomic absorption	NAA	Neutron Activation Analysis	XRF	X-ray fluorescence
β -c	Beta counting	PXE	PIXIE		
CAP	Capsule	SPEC	Spectrometer (SPEC = portable)		
Non-standard	AutoR	Auto regeneration	PTTL	Photo-transferred TL	

MINERALS & ETC.

cal	Calcite	Nf	Sodium feldspar	*	Other
ft	Flint	p	Polyminal	-	Not applicable
f	Feldspar	q	Quartz	e	Equivalent to (used as prefix)
Af	Unsep. alkali feldspar	z	Zircon	a	Year
Kf	Potassium feldspar	por	Porcelain		
Terms: I, P, a, b, A, S_N , S_O , TAC: as defined in the literature.					

Laboratory: Oxford	Entry: 31
--------------------	-----------

Site: Orp-le-grand Location: Brabant, Belgium Grid Ref.: 50° 41'40"N 4° 58'23"E.
 Site Description: Open air site, buried in upper part of loess in plateau situation.

Dates	Lab. Ref.	Material	Archaeological Reference
TL Single Ages: 13.7 ±1.7 ka (Ox88TLfg)	245a3	burnt flint	OP145/79
13.1 ±1.4	245a5		OP1164/26
12.9 ±1.5	245b1		OP1221/48
12.1 ±1.3	245b2		OP1133/79
11.8 ±1.2	245b4		OP1185/29

Archaeological Evidence: Magdalenian; as the chronological succession of the different Magdalenian groups in Western Europe is poorly understood these dates can be accepted as a good indication of the age of the Magdalenian of Orp.

Site Director: Prof. Dr P.M. Vermeersch. Catholic University of Leuven, Redingenstraat, 16 bis, 3000 Leuven, Belgium.

Reports: Vermeersch, P.M., Symens, N., Vynckier, P., Gijssels, G., Lauwers, R. (1987) Orp, site Magdalenien de plein air (comm.de Orp Jauche). *Archaeologia Belgica*, 111, 7-56.

PART II
 TECHNICAL SPECIFICATION

Section A. TL Measurements									
1. Min(fit) tech.(fig. 1 - 8µm)									
Sample Ref.	P ± s.e. (Gy)	I/P	Slopes	Plateau	Peak	Stability	a val.		
245a3	22.0 ± 1.5	0	-	± 3%; 300-350°	350°; 5°/s;	300 - 350°; 0.5 a; 18°; 100 ± 3%	0.09		
245a5	22.5 ± 1.5	0	-	± 3%; 325-375°	350°; 5°/s;	325 - 375°;	0.08		
245b1	19.7 ± 1.3	0	-	± 5%; 350-425°	375°; 5°/s;	350 - 425°;	0.06		
245b2	22.6 ± 1.5	0	-	± 3%; 300-375°	350°; 5°/s;	300 - 375°;	0.07		
245b4	20.8 ± 1.3	0	-	± 3%; 300-375°	350°; 5°/s;	300 - 375°;	0.05		

Section B. Dose-rate Measurements									
Sample Ref.	Total Eff. Dose-rate	Dose-rate Components			Radon		Water		
		α	β	γ	cos.	%	Sample	Env.	%
	mGy/a	%	%	%	%	%	%	%	%
245a3	1.60 ± 0.10	12	9	71	8	0 ± 5	0 ± 2	12 ± 3	-
245a5	1.71 ± 0.11	12	13	66	9	-	-	-	-
245b1	1.53 ± 0.10	8	12	71	9	-	-	-	-
245b2	1.86 ± 0.11	16	19	58	7	-	-	-	-
245b4	1.76 ± 0.11	13	18	61	8	-	-	-	-
Method		α-c	α-c	CAP	SpEC	α-c			
		FFH	FFH	SPFC					

Section C. Error [eA76]				
Section D. TL Age				
Sample Ref.	TL Age ka	Random ka	Errors ka	Overall ka
245a3	13.7	-	-	1.7
245a5	13.1	-	-	1.4
245b1	12.9	-	-	1.5
245b2	12.1	-	-	1.3
245b4	11.8	-	-	1.2

Laboratory: Oxford	Entry: 32
--------------------	-----------

Site: Nine Mile Water Location: Nether Wallop, Hampshire, UK. Grid Ref.: SU 307345
 Site Description: Raised mound adjoining Wallop Brook, on westerly edge of water meadow system.

Dates	Lab. Ref.	Material	Archaeological Reference
TL Context Age: 4240 ±380 a (Ox88TLfg)	260f	burnt flint	TL Pit layer 2
TL Context Comps:	4050 ±400		260f1
	4350 ±430		260f3
	4190 ±410		260f6
	4230 ±420		260f9
	4380 ±430		260f13

Archaeological Evidence: Excavation revealed a sequence of topsoil overlying burnt flint sealing a fine grey/white deposit. Pottery at the base of the topsoil has been dated to the 12th/13th Centuries. There are however some L.A. sherds and the extent of burnt flint is unknown. A pre-L.A. date is therefore expected. The molluscan analysis needs to be completed before further comment can be made.

Site Director: Diane Williams, Department of Archaeology, University College, P.O. Box 78, Cardiff CF1 1XL.
 Reports: D. Williams, in preparation.

PART II
 TECHNICAL SPECIFICATION

Section A. TL Measurements									
1. Min(fit) tech.(fig. 1 - 8µm)									
Sample Ref.	P ± s.e. (Gy)	I/P	Slopes	Plateau	Peak	Stability	a val.		
260f1	2.55 ± 0.20	0	-	± 5%; 325-375°	350°; 5°/s;	325 - 375°; 0.5 a; 18°; 100 ± 3%	0.13		
260f3	3.92 ± 0.35	0	-	± 3%; 350-400°	-	350 - 400°;	0.20		
260f6	2.60 ± 0.20	0	-	± 5%; 325-400°	-	325 - 400°;	0.11		
260f9	2.75 ± 0.20	0	-	± 3%; 325-400°	-	325 - 400°;	0.06		
260f13	3.02 ± 0.20	0	-	± 5%; 350-425°	-	350 - 425°;	0.11		

Section B. Dose-rate Measurements									
Sample Ref.	Total Eff. Dose-rate	Dose-rate Components			Radon		Water		
		α	β	γ	cos.	%	Sample	Env.	%
	mGy/a	%	%	%	%	%	%	%	%
260f1	0.63 ± 0.06	14	18	46	22	0 ± 3	0 ± 2	20 ± 5	-
260f3	0.90 ± 0.08	38	14	33	15	-	-	-	-
260f6	0.62 ± 0.06	18	13	47	22	-	-	-	-
260f9	0.65 ± 0.06	12	22	45	21	-	-	-	-
260f13	0.69 ± 0.06	16	22	42	20	-	-	-	-
Method		α-c	α-c	CAP	SpEC	α-c			
		FFH	FFH	SPFC					

Section C. Error [eA76]				
Section D. TL Age				
Sample Ref.	TL Age a	Random a	Errors a	Overall a
260f3	4050	-	-	400
260f6	4350	-	-	430
260f9	4190	-	-	410
260f13	4230	-	-	420
	4380	-	-	430

Site: Jels 2
Location: The site is at 'Jels-lakes' near the town of Jels in the southern part of Jutland, Denmark.
Grid Ref.: M3608
Site Description: A small hunters camp belonging to the Hamburgian Culture.

Dates	Lab. Ref.	Material	Archaeological Reference
TL Context Age:	10.2	±0.9 ka	(Ox88TLfg) HAM 1610 Oksensvad sogn 204
TL Context Comps:	9.4	±1.1	600a1 2203
	10.8	±1.0	600a3 2287

Archaeological Evidence: A very artefact-rich hunters camp belonging to the Hamburgian Culture. (12-13000 B.P.) A few Mesolithic artefacts were found within the excavation area. The TL date is not in agreement with the archaeological dating.

Site Director: Jorgen Holm, Haderslev Museum, Dalgade 7, 6100 Haderslev, Denmark.
Flemming Rieck, Skibshistorisk Laboratorium, Frederiksborgvej 63, 4000 Roskilde, Denmark.
Reports: Holm J., and Rieck, F. (1987) Die Hamburger Kultur in Dänemark, *Archaeologisches Korrespondenzblatt*, 17, 151-165.

PART II
TECHNICAL SPECIFICATION

Section A. TL Measurements									
1. Min(fit) tech.(fig : 1 - 8µm)									
Sample Ref.	P ± s.e. (Gy)	I/P	Slps	s Plateau	Peak	Stability	a val.		
600a1	4.25 ± 0.50	0	-	± 5%; 325-400°	375°; 5°/s;	325-400°; 0.5 a; 18°; 100 ±3%	0.15		
600a3	5.50 ± 0.50	0	-	± 5%; 325-400°	375°; 5°/s;	325-400°;	0.04		

Section B. Dose-rate Measurements												
Sample Ref.	Total Eff. Dose-rate mGy/a	Dose-rate Components					Radon		Water		Env.	
		α	β	γ	cos.	%	%	%	Sample %	%	%	%
600a1	0.45 ± 0.07	10	10	51	29	0 ± 2	0 ± 5	18 ± 6	-	-	-	-
600a3	0.51 ± 0.07	10	20	45	25	-	-	-	-	-	-	-
Method	α-c α-c α-c SPEC SPEC FPH											

Section C. Error [eA76]

Section D. TL Age			
Sample Ref.	TL Age ka	Random ka	Overall ka.
600a1	9.4	-	1.1
600a3	10.8	-	1.0

Site: Jels 1
Location: The site is at 'Jels-lakes' near the town of Jels in the southern part of Jutland, Denmark.
Grid Ref.: M 3608
Site Description: A small hunters camp belonging to the Hamburgian Culture.

Dates	Lab. Ref.	Material	Archaeological Reference
TL Context Age:	12.40	±1.60 ka	(Ox84TLfg) 600g(25) burnt flint HAM 1356
TL Context Comps:	12.85	±1.95	600g(25)ii 47.2
	12.10	±1.80	600g(25)iii 1625
	12.25	±1.80	600g(25)iii 1686

Archaeological Evidence: In N.Germany similar sites are dated by C-14 to the Bolling interstadial, ie 12-13000 B.P.

Site Director: Jorgen Holm, Haderslev Museum, Dalgade 7, 6100 Haderslev, Denmark.
Flemming Rieck, Skibshistorisk Laboratorium, Frederiksborgvej 63, 4000 Roskilde, Denmark.
Reports: Holm J., and Rieck, F. (1987) Die Hamburger Kultur in Dänemark, *Archaeologisches Korrespondenzblatt*, 17, 151-165.

PART II
TECHNICAL SPECIFICATION

Section A. TL Measurements									
1. Min(fit) tech.(fig : 1 - 8µm)									
Sample Ref.	P ± s.e. (Gy)	I/P	Slps	s Plateau	Peak	Stability	a val.		
g(25)ii	8.7 ± 0.5	0	-	± 5%; 325-400°	350°; 5°/s;	325 - 400°; 0.5 a; 18°; 100 ±3%	0.030		
g(25)iii	9.6 ± 0.5	0	-	± 5%; 325-400°	375°; 5°/s;	325 - 400°; 0.5 a; "	0.095		
g(25)iii	9.7 ± 0.5	0	-	± 5%; 325-400°	375°; 5°/s;	325 - 400°; 0.5 a; "	0.095		

Section B. Dose-rate Measurements												
Sample Ref.	Total Eff. Dose-rate mGy/a	Dose-rate Components					Radon		Water		Env.	
		α	β	γ	cos.	%	%	%	Sample %	%	%	%
g(25)ii	0.68 ± 0.08	4	13	65	18	0 ± 2	-	-	-	-	-	-
g(25)iii	0.80 ± 0.10	7	23	55	15	-	-	-	-	-	-	-
g(25)iii	0.80 ± 0.10	7	23	55	15	-	-	-	-	-	-	-
Method	α-c α-c α-c SPEC SPEC FPH											

Section C. Error [eA76]

Section D. TL Age			
Sample Ref.	TL Age ka	Random ka	Overall ka.
g(25)ii	12.85	-	1.95
g(25)iii	12.10	-	1.80
g(25)iii	12.25	-	1.80

Special Remarks:
• Environmental dose-rate were determined by V. Mejdahl, Risø National Laboratory using a portable gamma-ray spectrometer.

Site: West Overton
Location: River valley bottom in Upper Kennel, North Farm, West Overton, Wiltshire., UK.
Grid Ref.: SU/135686
Site Description: Cremation hearth.

Dates	Lab. Ref.	Material	Archaeological Reference
TL Context Age:	3030 ± 250 a	(Ox88TLfg)	727f burnt flint Long Meadow Pit D
TL Context Comps:	3090 ± 275 2870 ± 270 3130 ± 280	727(I) 727(II) 727(IV)	

Archaeological evidence: Bronze age (probably Deverel-Rimbury) pot, up-turned over a human cremation. It was in a buried soil covered by alluvium. The charcoal in the pot is C-14 dated to 3020 ± 70 BP (OxA-1348).

Site Director: J. G. Evans, Department of Archaeology, University College, PO Box 78, Cardiff CF1 1XL, UK.
Reports: For related C-14 date see *Archaeometry*, 29(2), 1987 pp 294-295 (Also forthcoming date list of AMS lab).

PART II
TECHNICAL SPECIFICATION

Section A. TL Measurements							
1. Min(f) tech.(fg ; 1 - 8µm)							
Sample Ref.	P ± s.e. (Gy)	I/P	Slps	Plateau	Peak	Stability	a val.
727(I)	2.38 ± 0.30	0	-	± 3%; 325-400°	375°; 5°%; -	325 - 400°; 0.5a; 18°; 100 ± 3%	0.10
727(II)	2.38 ± 0.30	0	-	± 3%; 325-375°	350°; 5°%; -	325 - 375°; "	0.08
727(IV)	2.38 ± 0.30	0	-	± 3%; 325-375°	350°; 5°%; -	325 - 375°; "	0.10

Section B. Dose-rate Measurements									
Sample Ref.	Total Eff. Dose-rate	Dose-rate Components				Radon	Water Sample	Env.	
		α	β	γ	cos.				
	mGy/a	%	%	%	%	%	%	%	%
727(I)	0.77 ± 0.05	16	12	55	17	0 ± 3	0 ± 2	32 ± 3	
727(II)	0.83 ± 0.05	16	17	51	16	"	"	"	
727(IV)	0.76 ± 0.05	14	12	56	18	"	"	"	
Method	α-c	α-c	α-c	CAP	SpEC	α-c			
		FPh	FPh	SpEC	SpEC				

Section C. Error [eA76]

Section D. TL Age			Errors	
Sample Ref.	TL Age a	Random a	Overall a	
727(I)	3090	-	275	
727(II)	2870	-	270	
727(IV)	3130	-	280	

Site: Bagaggera
Location: Merate village, Lecco region, Lombardy, N. Italy.
Grid Ref.: F 32 Como. 45°42'54" N., 9°23'18" E. 260 m above sea level.
Site Description: Pleistocene terrace.

Dates	Lab. Ref.	Material	Archaeological Reference
TL Single Age:	60.5 ± 7.5 ka	(Ox89TLfg)	750f burnt flint 282

Archaeological Evidence: Burned flint from a Mousterian level at the base of an Upper Pleistocene loess, 40 m thick; at the top of weathered fluvial Middle Pleistocene sand and gravel.

Site Director: Prof. Mauro Cremaschi, Dipartimento di Scienze della Terra, Via Mangiagalli 20133 Milano, Italy.
Reports: Preliminary reports in:
Cremaschi M., Orombelli G., Salloway J.C. (1985) Quaternary stratigraphy and soil development at the S. border of the Central Alps (Italy): the Bagaggera sequence. *Riv. It. Paleont. Strai.*, 90(4), 565 - 603.
Cremaschi M. (1987) Paleosols and Venusols in the central Po Plain (N. Italy). Thesis, University of Amsterdam, Unicopli, Milano, pp1-360. Definitive report in preparation.

PART II
TECHNICAL SPECIFICATION

Section A. TL Measurements							
1. Min(f) tech.(1 - 8 µm)							
Sample Ref.	P ± s.e. (Gy)	I/P	Slps	Plateau	Peak	Stability	a val
750f	99.8 ± 5.0	0	-	± 3%; 375-450°	375°; 5°%; -	375-400°; 0.5a; 18°; 100 ± 3%	0.10

Section B. Dose-rate Measurements									
Sample Ref.	Total Eff. Dose-rate	Dose-rate Components				Radon	Water Sample	Env.	
		α	β	γ	cos.				
	mGy/a	%	%	%	%	%	%	%	%
750f	1.65 ± 0.16	7	11	75	7	0 ± 5	0 ± 2	28 ± 7	
Method	α-c	α-c	α-c	SPEC	SpEC	α-c			
		FPh	FPh	SpEC	SpEC				

Section C. Error [eA76]

Section D. TL Age			Errors	
Sample Ref.	TL Age ka	Random ka	Overall ka	
750f	60.5	-	7.5	

Site: Three Ways Wharf.
Location: 101-105 Oxford Road, Uxbridge, Middlesex, UK.
Grid Ref.: TQ 052 846
Site Description: Multiperiod site composed of a Late Glacial flint and bone scatter and an early Mesolithic flint and bone scatter.

Dates	Lab. Ref.	Material	Archaeological Reference
TL Single Age: 8000 ± 800 a	(Ox89TLfg)	772fl burnt flint	UX88VIII E23/NO2/D/3 343 SF64

Archaeological Evidence: Scatter A contained c.700 pieces of flint with some affinity to the 'Long Blade' late glacial period, with associated fauna of reindeer and horse. This latter dated by C-14 to 10270 ± 100 B.P. (OxA1778) and 10010 ± 120 B.P. (OxA1902). Scatter C produced c.7000 pieces of struck flint and c.2000 fragments of bone. The lithic assemblage is early Mesolithic and fauna is dominated by red deer. The dated sample is from this assemblage contained within argillaceous sediments deposited by gentle overbank flooding. The flint work is dated typologically to the early Mesolithic (expected to date to ca 9000 BP). *JSCl*

Site Director: John S.C. Lewis, Dept. of Greater London Archaeology, Museum of London, Town Mission Hall, Mission Square, Pottery Road, Brentford, Middlesex TW8 0SD, UK.

Reports: Lewis, J. S. C. (1989) Excavations at Three Ways Wharf. *Mesolithic Miscellany*, 10.
Lewis, J. S. C. (1990) Excavations at Three Ways Wharf. Proc. The Late Glacial of N.W. Europe' (ed N. Barton), Dept. External Studies, Oxford University September 1989. In press.

PART II
TECHNICAL SPECIFICATION

Section A. TL Measurements						
1. Min(t) tech. (fig. 1 - 8µm)						
Sample Ref.	P ± s.e. (Gy)	I/P	Slps	Plateau	Peak	Stability
772fl	9.6 ± 0.6	0	-	± 5%; 325-400°	375°; 5°/s;	325-400°; 0.5 ± 18°; 100 ± 3%
						a val
						0.10

Section B. Dose-rate Measurements											
Sample Ref.	Total Eff. Dose-rate	Dose-rate Components						Radon		Water	
		α	β	γ	cos.	%	%	%	%	Sample	Env.
772fl	1.2 ± 0.2	10	8	71	11	0 ± 5	0 ± 2	0 ± 2	30 ± 8		
Method		α-c	α-c	SPEC	SPEC	FPPh	α-c				

Section C. Error [eA76]

Section D. TL Age		
Sample Ref.	TL Age ka	Errors Random ka Overall ka
772fl	8.0	- 0.8

Site: Bir Sahara, BS-1
Location: Bir Sahara East Depression in south western Egypt.
Grid Ref.: 22°52' N, 28°50' E
Site Description: Middle Palaeolithic. Samples are from buried layers above an occupation horizon.

Dates	Lab. Ref.	Material	Archaeological Reference
TL Single Ages: 105.4 ± 10.5 ka 108.6 ± 10.6 ka	(Ox88TLqi) (Ox88TLqi)	506a burnt 506b quartz	BS1TR965, V.H.73 upper layer lower layer

Archaeological Evidence: Large concentration of artefacts (approx. 230 per m²) of Levallois technique core and flakes; also highly denudate tool assemblage.

Site Director: Fred Wendorf, Dept of Anthropology, Southern Methodist University, Dallas, Texas 75275, USA.

Reports: Wendorf, F., and Schild, R. (1980) *Prehistory of the Eastern Sahara*. Academic Press, New York.

PART II
TECHNICAL SPECIFICATION

Section A. TL Measurements						
1. Min(q) tech. (inc; 90 - 150µm)						
Sample Ref.	P ± s.e. (Gy)	I/P	Slps	Plateau	Peak	Stability
506a	310 ± 30	0	1.05 ± 0.05	± 3%; 325-375°	350°; 5°/s;	-
506b	304 ± 30	0	1.05 ± 0.05			-

Section B. Dose-rate Measurements											
Sample Ref.	Total Eff. Dose-rate	Dose-rate Components						Radon		Water	
		α	β	γ	cos.	%	%	%	%	Sample	Env.
506a	2.94 ± 0.44	-	62	34	4	0 ± 5%	5 ± 5	7 ± 5			
506b	2.80 ± 0.42	-	57	39	4						
Method		α-c	α-c	SPEC	SPEC	FPPh	α-c				

Section C. Error [eA76]

Section D. TL Age		
Sample Ref.	TL Age a	Errors Random a Overall a
506a	105.4	- 10.5
506b	108.6	- 10.6