

www.ancienttl.org · ISSN: 2693-0935

Issue 19(1) - June 2001 https://doi.org/10.26034/la.atl.v19.i1

This issue is published under a Creative Commons Attribution 4.0 International (CC BY): https://creativecommons.org/licenses/by/4.0



© Ancient TL, 2001

Manual coring equipment for the collection of stratified samples from dry sand dunes

Olaf Bubenzer

University of Cologne, Department of Geography, Albertus-Magnus-Platz, D-50923 Köln, Germany, e-mail: <u>olaf.bubenzer@uni-koeln.de</u>, Fax.: 0049-(0)-221-470-5124, Tel.: 0049-(0)-221-470-3966

(Received in final form 8 January 2001)

Introduction

The analysis of the sandy layers of dunes, especially from draas (or megadunes), are suitable for dating palaeo-processes and the reconstruction of palaeo-windsystems (Besler, 1998). The sand layers in dunes are created by different wind directions and/or velocities and often show different grain sizes and colours. Thus, many dunes show a clear structure and their separate analysis is important for a large of sedimentological, geomorphological, range climatological and archaeological studies. Dry, nonfixed sand with no observable cohesion or pedogenetic developments behave in some way similar to liquids (Haynes and Johnson, 1984). Therefore, it is difficult to collect samples by drilling or digging because the sand flows immediately back into the hole. Until now it was only possible to reach depths of 1-2 m with manual drilling methods (Besler and Gut, 1997). Drilling to greater depths required heavy machines and large water supplies, which is not feasible for sampling in remote areas. Other problems arise from the fact that samples for luminescence dating have to be undisturbed and not be exposed to light. Furthermore, sample sizes in the range of about 500g are required. Wallinga and van der Staay (1999) described a suction corer, which is suitable for collecting samples from waterlogged sands. This paper describes a lightweight manual coring equipment for dry sands to reach deep sediment layers and to collect undisturbed samples. It was developed within the Collaborative Research Center ACACIA (Arid Climate, Adaptation and Cultural Innovation in Africa, SFB 389, University of Cologne) - section A "Holocene Environment and Cultural History in Northeastern Africa.

Equipment, drilling and sampling method

Figure 1 shows a schematic drawing of the drilling and sampling equipment, which can be used for sample collection to depths of up to 6 m. It weighs about 25 kg for the basic tools and the sampling equipment and 6 kg/m for the tubing. It is available from Eijkelkamp Co., P.O. Box 4, 6987 ZG Giesbeek, The Netherlands (www.eijkelkamp.com). The water consumption is about 1-2 l/m.

The coring equipment can be used in two configurations:

(i) Collection of mixed samples

The plastic liner tube (1) is placed on the sand surface. A light drilling-rod with a bayonet connection (2) and an Edelman sand auger (3) is inserted and used to core below the liner. Movable casing tube clamps (4) are used to hold the liner in place and to press it down whilst lifting the auger. For the first 1 to 2 m no water supply is necessary because it takes only a very short time to lift the auger. At greater depths, about 200 ml water is needed for one auger scoop of 10 to 20 cm. The full auger produces about 250 cm³ sand. The coring works by cutting, therefore, the auger samples are mixed over the average coring depth of 10-20 cm.

(ii) Collection of undisturbed, stratified samples

After the liner has been driven to a given depth, it is possible to collect stratified samples with diameters of 6.5 cm and lengths of 40 cm. A removable stainless steel tube (6) with a core catcher (7) on the head of a rod with a conical



Fig. 1: Schematic drawing of the coring equipment

screw thread connection (5) is pushed with a percussion-free hammer into the sediment, which is watered prior to the insertion of the equipment. The watering in conjunction with core catcher ensure that the sediment is hauled as a complete sample core. The inner PVC-foil ensured that the sediment cores stays intact. The stainless steel tubes can be disconnected from the rod and taken to the laboratory where the sample is extracted in darkroom conditions. The maximum sediment volume in a sample tube is 1.400 cm^3 . Two persons will require about 2 to 3 hours to reach a depth of 3 m and collect one sample.

Depending on grain size and sediment density it is possible - in our experience - to reach maximum depths of about 6 m. The equipment is maintenancefree and can easily be carried a few hundred meters without a car.

References

- Besler, H. (1998) Periods of aeolian activity and stability in the Great Sand Sea in Egypt. Zbl. Geol. Paläontol, I, 1997, 23-39.
- Besler, H. and Gut, S. (1997) Untersuchungen zum Feuchtegehalt in D
 ünen am Beispiel der Namib/Namibia. Kölner Geogr. Arb., So.-Folge Afrika, 13, 39-47.
- Haynes, V.C. and Johnson, D.L. (1984) A provisional soil chronosequences of the Quaternary of the eastern Sahara. *Geological Society of America with Programs*, 16, 534-535.
- Wallinga, J. and van der Staay, J. (1999) Sampling in waterlogged sands with a simple handoperated corer. Ancient TL, 17, 59-61.

Reviewer Rainer Grün

Dose evaluation on a fossil tooth using multiple and single powder aliquots as well as a fragment

Rainer Grün, Steve Robertson

Research School of Earth Sciences The Australian National University Canberra ACT 0200, Australia

Matthew Gordon Daramalan College Canberra ACT 2602

(Received 12 February 2001; in final form 28 February 2001)

Introduction

The use of single aliquots for ESR dating of tooth enamel has the advantage that small samples can be analysed (5 to 30 mg), requiring less sample preparation time, and that sample inhomogeneity becomes insignificant. Single aliquots dose measurements can be carried out on powders as well as fragments. The analysis of the latter is necessary when working on valuable samples, such as fossil human teeth. Thus far, no experiments have been carried out to validate single aliquot measurements on powders nor fragments.

Experimental

We have selected sample 698, a relatively large bovid molar for the archaeological site of La Quina, which allowed the preparation of a series of sub-samples. The following procedures for dose estimation were tested (for the relative location of the sub-samples see Figure 1):

- conventional multiple aliquots, one set (M2) heated for 2h at 90 °C after irradiation, then stored for at least 1 day at ambient temperatures and humidity following the recommendations of Hayes et al. (2000), the other set (M1) not heat treated;
- single powder aliquots, not heated treated (S1 to S4);
- single powder aliquots, heated for 2h at 90 °C and equilibrated after each irradiation step (H1 to H4);
- 4) a fragment, not heat treated (F2 was measured only).

The enamel of the fragments and single aliquots had an average thickness of about 1200 μ m, whilst the enamel of the multiple aliquots was thinner (800 to $1000 \ \mu m$).



Figure 1.

Schematic drawing of the positions of the samples. M: multiple aliquots, S: single aliquots, not preheated; H: single aliquots, preheated. The enamel in the area of M2 was thinner than for the other samples.

Samples were irradiated with 0, 10.2 20.4, 40.8, 81.6, 132.6, 163.2 and 204 Gy using a 137 Cs source and measured using routine measurement conditions. The fragment was mounted in a quartz holder such that the outer and inner surfaces of F2 were perpendicular to the rotational axis. The sample was measured in 10° intervals with a programmable goniometer. For spectrum evaluation of the powders, we used a series of well described procedures (Grün 2000), plus natural spectrum fitting (scaling the



Figure 2.

(A) Dose values for each sample using a variety of evaluation methods
(B) Averaged dose values for the sample groups for a variety of evaluation methods
(C) Average, weight normalised ESR intensities of the single aliquots, non-heated and heated. The data points are slightly offset for clarity

natural spectrum into the irradiated spectra, for more details see Grün, submitted). The latter study concluded that natural spectrum fitting ought be the most reliable method for spectrum evaluation because it was least dependent on interferences. For the fragment, spectrum deconvolution (using four Gaussian peaks: see Robertson and Grün 2000) as well as natural spectrum fitting was applied. The dose values were obtained by linearisation of the single saturating exponential dose response function and Monte Carlo error calculation (Grün and Brumby 1994).

Results and Discussion

Figure 2 presents the results on the powder samples. The dose results for each sub-sample are not dependent on the method of spectrum evaluation (Figure 2A): the dose values of a given sub-sample agree within error (there may be some tendency for lower dose estimation when using integral deconvolution for samples S1, S2 and S4). The dose values within each group of samples (i.e. multiple aliquots, single aliquots: non heated, single aliquots: heated) also agree within error. There is also no quantifiable difference between M1 (not heated) and M2 where each aliquot was heated after irradiation.

Figure 2B shows the average dose values for each sample group. Again, the dose values show little dependence on the method of spectrum evaluation. The dose results of the multiple aliquots and the single, non-heated aliquots agree within error. The small systematic offset between the means of the multiple aliquots and the non-heated single aliquots can be attributed to the difference in enamel thickness. However, it seems clear that repeated preheating leads to significant smaller doses when using single aliquots (Figure 2B). The reason for the dose differences lie in changes in the natural, as well as the dosed samples. The heating decreases the natural intensity by about 9% whereas the intensity of the highest dose point increases by about 11% (Figure 2C). The effect could be explained by the presence of a less stable component (or indeed part of the stable component, depending on the kinetics) which is removed from the natural sample and that some other charges (not present in the natural sample) are re-distributed after irradiation (see also Brik et al. 2000, Hayes and Haskell 2000). Heating experiments by Scherbina and Brik (2000) indicate

some charge transfer between the CO_2 radicals with other "unobservable" centres at temperatures above 80°C. Alternatively, the heating could lead to the generation of additional precursors similar to those observed in speleothems (Yokoyama et al. 1985), mollusc shells (Brumby and Yoshida 1994) and corals (Yoshida and Brumby 1999)

Unfortunately, spectrum analysis does not show an unambiguously identifiable component that is apparently re-distributed in the spectra. However, the results clearly demonstrate that the prescribed heating procedure does not simulate geological aging, because the overall effect on the irradiated spectra ought be qualitatively the same as for the natural spectra (i.e. an overall decrease in the ESR intensity should occur). It should be noted here that experiments by Skinner et al. (2000) showed an increase of the dose value of a non-heated sample over 6 months. They concluded that this process could be simulated by preheating the aliquots for 3 days at 90 °C. On the other hand, Rhodes and Grün (1991) did not detect any changes in the dose values of two non-heated samples 2 years after the initial measurement. Whilst it is at present not possible to state that preheating is generally to be avoided, the setting chosen in this study is clearly inappropriate for single aliquots of geological samples. The experience from a large number of studies implies that ESR results obtained on non-heated samples does not yield systematic age overestimations. Indeed, samples from La Quina have caused consternation because they seem to yield ages, which are systematically too young (see Grün et al. 1997).

Figure 3 shows the powder spectra of M1 (Figure 3A) and the spectra for the enamel fragment F2 at 120° (Figure 3B) and at 220° (Figure 3C). These two angles show the largest spectral differences. Because of the angular distortions caused by the anisotropic crystal magnetic field of the hydroxyapatite fragment, the resulting spectra cannot be fitted with an ideal

powder CO_2 spectrum. The spectra of the fragment were analysed using spectrum deconvolution with four Gaussian lines (for more details see Robertson and Grün 2000), where Ax1 and Ax2 present the low and higher field component of the central, apparently axial signal. Peak-fitting with the natural spectrum was applied by optimising for the "whole peak" between S and E (see Figure 3A) as well as selected partial regions: T1 (between half height T1 and zero passing between T1 and B1), B1 (zero passing to half height between B1 and T2), B2 (half height between T2 and B2 and B2 to zero), the combinations of T1-B1, T1-B2 as well as T1-B2 width (S to T1 and B1 to E) and peak width (S to half height of T1 region and half height B2 to E). The latter two fitting methods are less dependent on interferences within the central peak region. Peak-to-peak methods were used for T1, B1, B2, T1-B1 and



Figure 3.

ESR spectra of a powder (A) and the fragment measured at angles of 120° (B) and 220° (C). The positions S, T1, B1, T2, B2 and E denote positions used for peak-to-peak measurements and to define regions for natural spectrum fitting.

T1-B2. The dose values of all spectrum evaluation methods show angular dependencies (Figure 4). At about 120°, the T1-B1 peak is largest and the dose values derived from Ax1, Ax1 + Ax2 as well as natural spectrum fitting are about the highest. At 220°, where B2 is largest, doses derived from Ax2 are highest, those of Ax1, Ax1 + Ax2 as well as natural spectrum fitting about lowest, respectively. Natural spectrum fitting yields average dose results that are close to the average dose of the single, nonheated aliquots, whereas all dose results from the deconvolution (Ax1+Ax2) lie below this value.

Figure 5 shows the average dose values for the various methods of spectrum evaluation for the fragment.. The errors bars represent the magnitude of dependency. angular the dose Spectrum deconvolution (Ax1 + Ax2) shows the least angular dependency, but also yields the smallest dose value as well as the largest deviation from the average dose value of the single aliquot powders (if sample selection and spectrum evaluation were ideal, the dose value of the fragment should be equal to the dose value of the single aliquot powder). All regions of interest of the natural fitting as well as all peak-topeak dose evaluations, albeit associated with larger errors, agree with the average dose value of the single aliquot powders. The reason why spectrum deconvolution of the absorption spectrum yielded the largest deviation from the average dose value of the powders most likely lies in the correct placement of the "wide" line in the deconvolution process (for more details see Robertson and Grün 2000).

When the dose response curves are normalised on the natural intensity, it can be observed that the angular dose dependency is caused by an angular sensitivity change. Similar observations were made by Brik et al. (2000) who attributed these interferences to a chaotic CO_2^- radical. Alternative explanations involve phase sensitive interferences or some anisotropic, non-radiation sensitive peaks. At the moment, however, we are not able to unambiguously identify these interferences or provide a plausible explanation for them.

Summary

Dose evaluations on multiple and single aliquots of powders yield consistent results. However, post irradiation heating causes severe dose underestimations. Although is it difficult to pinpoint the interferences which cause the intensity changes in the heated samples, it is clear that the prescribed heating does not simulate the geological ageing process, at least for this sample. We find that the average dose value of the single aliquot powders lie within the dose range provided from a variety of spectrum methods on the spectra of the fragment. Spectrum deconvolution, though being by far the most time intensive method, yielded the largest deviation between the dose values of the fragment and the powders, respectively.

It is clear that more experiments are required to pinpoint (and eliminate) any interfering signals in the ESR spectra of powders and fragments and that systematic fading tests ought be carried out. At seems at this stage, that pre-heating procedures should be treated with suspicion.

References

- Brik, A., Haskell, E., Brik, V., Scherbina, O. and Atamanenko, O. (2000) Anisotropy effects of EPR signals and mechanisms of mass transfer in tooth enamel and bones. *Applied Radiation* and Isotopes 52: 1077-1083.
- Brumby, S. and Yoshida, H. (1994) ESR dating of mollusc shells: investigations with modern shell of four species. Quaternary Geochronology 13: 157-162.
- Grün, R. (2000) Methods of dose determination using ESR spectra of tooth enamel. *Radiation Measurements* 32: 767-772.
- Grün, R. (submitted) ESR dose estimation on fossil tooth enamel by fitting the natural spectrum into the irradiated spectrum. *Radiation Measurements*.
- Grün, R., Abeyratne, M., Head, J., Tuniz, C. and Hedges, R.E.M. (1997) AMS ¹⁴C analysis of teeth from archaeological sites showing anomalous ESR dating results. *Quaternary Science Reviews* 16: 437-444.
- Grün, R. and Brumby, S. (1994) The assessment of errors in the past radiation doses extrapolated from ESR/TL dose response data. *Radiation Measurements* 23: 307-315.
- Hayes, R.B. and Haskell, E.H. (2000) A method for the identification of interfering signals in EPR dating of tooth enamel. *Radiation Measurements* 32: 781-785.
- Hayes, R.B., Haskell, E.H., Barrus, J.K., Kenner, G.H. and Romanyukha, A.A. (2000) Accurate EPR radiosensitivity calibration using small sample masses. Nuclear Instruments and Methods in Physical Research A 441: 535-550.
- Rhodes, E.J. and Grün, R. (1991) ESR behaviour of the paramagnetic centre at g=2.0018 in tooth enamel. Ancient TL 9: 14-18.





Figure 4.

(A) Dose values of the fragment F1 derived from spectrum deconvolution (for details see Robertson and Grün, 2000)

(B) Comparison of the dose values of the fragment obtained by spectrum deconvolution and natural spectrum fitting



Figure 5.

Summary of dose results. If there were no interferences, the dose of the fragment ought be the same as of the non-heated, single aliquots (dotted line). All evaluation strategies on the fragment yield dose values that are statistically indistinguishable from the single aliquots. There seems a tendency towards smaller dose values in the B2 region (see values for B2 region and B2 intensity).

- Robertson, S. and Grün, R. (2000) Dose determination on tooth enamel fragments from two human fossils. *Radiation Measurements* 32: 773-779.
- Scherbina, O.I. and Brik, A.B. (2000) Temperature stability of carbonate groups in tooth enamel. *Applied Radiation and Isotopes* 52: 1071-1075.
- Skinner, A.R., Blackwell, B.A.B., Chasteen, N.D., Shao, J. and Min, S.S. (2000) Improvements in dating tooth enamel by ESR. Applied Radiation and Isotopes 52: 1337-1344.
- Yoshida, H. and Brumby, S. (1999) Comparison of ESR ages of corals using different signals at X- and Q-band: re-examinations of corals from Huon Peninsula, Papua New Guinea. *Quaternary Science Reviews* 18: 1529-1536.
- Yokoyama, Y., Bibron, R., Leger, C. & Quaegebeur, J.P. (1985) ESR dating of paleolithic calcite: fundamental studies. *Nuclear Tracks and Radiation Measurements* 10: 929-936.

Reviewer

H. Schwarcz

Light transmittance through dry, sieved sand: some test results

Jeff Ollerhead

Department of Geography, Mount Allison University, 144 Main St., Sackville, New Brunswick, Canada E4L 1A7

(Received 19 February 2001; in final form 8 March 2001)

Abstract: a series of measurements was made at different light wavelengths of transmittance through various thicknesses of dry sand grains from a variety of size classes. Transmittance was found to decrease approximately exponentially with increasing sand layer thickness and with decreasing grain size. Transmittance did not vary significantly with incident light wavelength. It is concluded that for any dry sand, a layer with a thickness of 7 mm, and possibly much less, is capable of reducing the transmittance of incident light to 0.05 % or less. This result should be considered when collecting a surface sample as a modern analogue for a sample to be optically dated.

Introduction

It is now generally accepted that optical dating techniques can be used to determine the length of time that has elapsed since a mineral was last exposed to daylight (Aitken, 1998; Huntley and Lian, 1999). However, as Wintle and Huntley (1982) noted almost two decades ago, a necessary criterion for a valid luminescence age is that a modern sample in the same context yield an age of zero. Thus, in an 'ideal world', a modern sample (usually a surface sample) from an environment under study yields an optical age of zero. However, this is not always the case. For example, Wintle et al. (1994) obtained an optical age of 40 ± 15 years for a surface sand sample. In my own work, I have obtained optical ages significantly greater than zero for modern samples. An age greater than zero for a modern sample can occur because: (i) the sample is not well bleached prior to collection, (ii) there is a deficiency in the dating technique used (e.g., an inappropriate laboratory bleaching procedure, such as one which uses photons not present in nature, is used), or (iii) both (i) and (ii). The focus here is on one factor related to bleaching: sediment that might block overlying light penetration.

Modern samples should normally be collected as analogues for older samples being dated. If modern samples are found to be inadequately bleached, this raises questions about whether older samples of interest were adequately bleached prior to deposition. An understanding of the environment from which a sample is collected is clearly needed to address the question of whether or not one should expect mineral grains in a modern sample to be bleached sufficiently to yield an optical age of zero. In my own work, I have usually collected the top 0.5-1.0 cm of sediment from a site as a modern sample. This is done on the assumption that this layer of grains is well bleached. In an active environment (*e.g.*, on a beach) this is probably a fair assumption as the grains are usually in motion every few days or weeks. Nevertheless, it became apparent that a more concrete understanding of the magnitude of light attenuation through a given thickness of sediment is desirable.

A search of available literature revealed little data to aid in this understanding. Others have studied the bleaching process (e.g., Huntley, 1985; Berger, 1990; Gemmell, 1994; Huntley and Clague, 1996) and most review papers on the subject (e.g., Berger, 1995; Huntley and Lian, 1999) suggest various mechanisms for incomplete bleaching (e.g., overlying water, rapid burial, coatings on grains, etc.). However, the only paper found directly related to this topic was that of Ditlefsen (1992) who investigated the bleaching of K-feldspars in turbid water suspensions. His approach was to bleach samples in different suspensions and then measure their luminescence and compare it to that of unbleached grains. He showed that in dense suspensions (> 0.05 g/l) little bleaching took place. This led him to consider the limitations of using optical dating techniques to obtain ages for water-laid sediments.

From Ditlefsen's (1992) results one can infer that light attenuation through dry sediments should be very rapid. The purpose of this study was to test this idea. The key variables were assumed to be: (i) sediment thickness, (ii) grain size, and (iii) light wavelength. Transmission of light through sand can happen in two ways: (i) transmission through grains, and (ii) transmission through spaces between grains. It was assumed that most of the light would be transmitted through spaces between grains although this was not tested. It was expected that light transmission would decrease exponentially with sand thickness. It was also expected that light transmission would decrease with decreasing grain size because, although the pore volume would be unchanged, there would be more reflections at grain surfaces and during each reflection a significant fraction of the intensity is not reflected. Given that the sand tested contained different minerals (e.g., quartz, K-feldspar, etc.), a variation in light transmittance with wavelength was anticipated although the nature of such an effect was unknown. In order to test the hypotheses, a series of measurements were made at different light wavelengths of transmittance through various thicknesses of sand grains from a variety of size classes.

Sample collection, preparation and measurement

A large (~ 1 kg) sand sample was collected at Cape Jourimain, New Brunswick. In the laboratory it was rinsed with distilled water and then dried for 24 hours at 105°C. It was then sieved, in batches, to extract the following size fractions: -2.0, -1.0, -0.5, 0, 0.5, 1.0, 1.5, 2.0, 2.5, 3.0, 3.5 and 4.0 \$\phi\$. In each case the size fraction value is that of the smallest grains in a fraction (e.g, the 1.0 ϕ fraction contains grains that fit through a sieve with 0.5ϕ openings but not through a sieve with 1.0 ϕ openings). The mm equivalents for each ϕ size are provided in Table 1.

A custom holder was machined to hold a pair of standard microscope slides (25 mm x 75 mm x 1 mm thick) at the following spacings: 1.0, 1.5, 2.0, 2.5, 3.0, 4.0, 5.0, 7.0, 9.0, and 11.0 mm. To establish a reference prior to each measurement, transmittance for a given spacing with no sample was measured from 320 to 820 nm at 2 nm increments using a Hewlett Packard 8452A diode-array spectrophotometer. Sand grains from a given size class were then poured into the space between the two microscope slides and transmittance measured from 320 to 820 nm at 2 nm increments. The instrument used is a single-beam, microprocessor controlled, UV/VIS spectrophotometer with collimating optics. The beam is 6 mm x 8 mm and its intensity is approximately 28 mW·cm⁻². Wavelength accuracy is ± 2 nm and wavelength reproducibility is ± 0.05 nm. The lower limit of measurable transmittance is 0.05 %. Some grain-size classes could not be measured at some spacings because the

Sand thickness required Grain-size class to reduce T to ≤ 0.05 % (**þ**) (mm)(mm)-2.0 4.00 7 -1.0 2.00 5 -0.5 1.41 5 0 1.00 4 0.5 0.71 3 1.0 0.50 2 1.5 0.35 1.5 2.0 0.25 1 2.5 0.18 < 1 3.0 0.13 < 1 3.5 0.09 < 1 0.06 < 1

4.0

Table 1.

Thickness of sand required (in mm) to reduce the transmittance of incident light to 0.05 % or less (the lower limit of the instrument used is 0.05 %) for the grain-size classes tested for incident light wavelengths from 320 to 820 nm. Transmittance was found to be independent of wavelength so this variable is not included in the table.

Measurement results

Representative examples of the results obtained are presented in Figures 1 and 2. As expected, transmittance was found in all cases to decrease approximately exponentially with increasing sand layer thickness. For example, Figure 1(a) shows transmittance of about 7 % for a 1.5 mm thick layer of grains from the 1.4 mm size class; transmittance drops to about 1 % for a 3.0 mm thick layer of the same grains. Figure 1(b) shows that transmittance is only about 1 % for a 1.5 mm thick layer of grains from the 0.7 mm size class. It was also found that transmittance decreases approximately exponentially with decreasing grain size. For example, Figure 2 shows the rapid decrease in transmittance with decreasing grain size for a layer of sand 1 mm thick. As Figures 1 and 2 also indicate, no observable dependence on incident light wavelength was found.

grains were larger than the space between slides. All results presented are for the mean of two independent trials.



Figure 1.

Relationship between light transmittance and sand layer thickness for light at 400, 600, and 800 nm for the (a) 1.4 mm, and (b) 0.7 mm grain-size classes.

Table 1 shows the thickness of sand required (in mm) to reduce the transmittance of incident light to 0.05 % or less for the grain-size classes tested. These data show that a sand layer thickness of 7 mm is sufficient to reduce the transmittance of incident light to 0.05 % or less at all wavelengths tested for any grains smaller than 4 mm in diameter. They also show that a sand layer thickness of 1 mm is sufficient to reduce the transmittance of incident light to 0.05 % or less for all wavelengths tested for any grains smaller than 0.25 mm in diameter.





Relationship between light transmittance and grain size for light at 400, 600, and 800 nm for a layer of sand 1 mm thick.





Sand layer thickness required to reduce the transmittance of 600 nm incident light to 0.05 % or less (the lower limit of the instrument used is 0.05 %) for the 8 largest grain-size classes tested.

Figure 3 summarizes the results presented in Table 1 for 600 nm incident light. It shows the sand layer thickness required to reduce transmittance to 0.05 % or less for a given grain-size class. Although there is scatter in the data, primarily due to the fact that discrete sand layer thicknesses were used, the nature of the relationship is apparent.

Discussion and conclusions

The results show that light transmittance decreases approximately exponentially with increasing sand layer thickness and with decreasing grain size. The greatest amount of scatter in the data was for the larger grain sizes. This may be due to much greater variability in the packing of the larger grains which tend to be more angular than their smaller diameter counterparts. In supplemental testing, tapping the microscope slides to consolidate grains in the larger size classes (≥ 2 mm) resulted in reductions of transmittance by a factor of two or three. For grains from the smaller size classes (≤ 1 mm), tapping the microscope slides had no measurable effect on transmittance. Note that 'natural sands' are likely to be well packed and include a variety of grain sizes and thus transmittance through them is likely to be much less than what is reported here for sands with a very limited grain-size range.

No relationship between light transmittance and incident light wavelength was observed. This does not mean that a relationship does not exist but it does suggest that light wavelength is not an important variable in predicting rate of light attenuation through dry sand where more than a monolayer is present. It may also be an indication that, as assumed, most of the light is being transmitted around the grains rather than through them.

The results indicate that for any sand, a layer with a thickness of 7 mm, and possibly much less, is capable of reducing the transmittance of incident light to 0.05 % or less. This should be taken into account when collecting a surface sample as a modern analogue for a sample to be optically dated. As suggested previously, collecting the top 10 mm of a sand in an active environment is probably reasonable since one would expect most of the grains to be in motion every few days or weeks and thus be well bleached. In a less active environment (e.g., the bed of a sheltered lake), making this assumption is probably not wise. One should try to collect only the actual surface grains as a modern analogue in this case. One technique for doing this is by using adhesive tape as first described by Readhead (1984).

Acknowledgements

Lynn Borse (field assistance), Karen Sekikawa and Marie Tower (laboratory assistance), Al MacDonald (machining assistance), and John Read are thanked for helping with this project. David Huntley, Jack Rink and John Prescott are thanked for their advice and for commenting on drafts of this paper. Financial support was provided by the Natural Sciences and Engineering Research Council of Canada.

References

- Aitken, M.J. (1998). *Introduction to optical dating*. Oxford University Press, Oxford.
- Berger, G.W. (1990). Effectiveness of natural zeroing of the thermoluminescence in sediments. *Journal of Geophysical Research*, **95**, 12375-12397.
- Berger, G.W. (1995). Progress in luminescence dating methods for quaternary sediments. *In*: Rutter, N.W. and Catto, N.R. (eds), *Dating methods for Quaternary deposits*, pp. 81-104. Geotext2, Geological Association of Canada, St. John's.
- Ditlefsen, C. (1992). Bleaching of K-feldspars in turbid water suspensions: a comparison of photo- and thermoluminescence signals. *Quaternary Science Reviews*, **11**, 33-38.
- Gemmell, A.M.D. (1994). Environmental controls on the TL age of modern (zero-age) proglacial outwash sediments. *Quaternary Geochronology (QSR)*, **13**, 485-489.
- Huntley, D.J. (1985). On the zeroing of the thermoluminescence of sediments. *Physics and Chemistry of Minerals*, **12**, 122-127.
- Huntley, D.J. and Clague, J.J. (1996). Optical dating of tsunami-laid sands. *Quaternary Research*, 46, 127-140.
- Huntley, D.J. and Lian, O.B. (1999). Using optical dating to determine when a sediment was last exposed to daylight. *In*: Lemmen, D.S. and Vance, R.E. (eds), *Holocene climate and environmental change in the Palliser Triangle: A geoscientific context for evaluating the impacts of climate change on the southern Canadian prairies*, pp. 211-222. Geological Survey of Canada Bulletin 534, Ottawa.
- Readhead, M.L. (1984). *Thermoluminescence dating* of some Australian sedimentary deposits. PhD thesis, Australian National University, Canberra.
- Wintle, A.G. and Huntley, D.J. (1982). Thermoluminescence dating of sediments - A review. *Quaternary Science Reviews*, **1**, 31-53.
- Wintle, A.G., Lancaster, N. and Edwards, S.R. (1994). Infrared stimulated luminescence (IRSL) dating of late-Holocene aeolian sands in the Mojave Desert, California, USA. *The Holocene*, **4**, 74-78.

Reviewer

J.R. Prescott

Comments

This is a useful piece of laboratory work in support of field measurements and is helpful in interpreting them. It is surprising how little direct laboratory work of this kind has been reported. Since it also good physics, it appeals to me. My generation would probably ask, "What did Rayleigh say about it?" The exponential absorption relation and its eventual breakdown would come as no surprise to him. For clean quartz, no wavelength dependence is expected for particles in this size range. In the wider luminescence dating world, almost everyone one who has tried it, finds that the surface layer 0.5-1.0 cm has a small but non-zero age, and/or that quartz bleached in layers in the laboratory reaches a lower level than surface quartz collected in the field. The present paper throws some light on this, if the play on words may be forgiven.

Such measurements were carried out quite early in the application of luminescence techniques to sediments. Ollerhead refers to Readhead's thesis (1984). This relatively inaccessible reference contains an account of considerable laboratory work, which was also reported, in part, at the Cambridge LED meeting: Readhead, Quater. Sci. Rev. 7, 257-264 (1988). Historians may also care to look up the Helsingor LED Conference Proceedings at: Council of Europe Journal PACT 9, 505-512 (1983). Modesty forbids naming the author.

It is worth adding that there is a difference between TL and OSL in that some samples are never completely zeroed for TL, no matter how long they are exposed to sunlight.

Comparison of dose rate determination using high – resolution gamma spectrometry and inductively coupled plasma – mass spectrometry

Frank Preusser^{1,2} and Haino Uwe Kasper¹

¹Geologisches Institut, Universität zu Köln, Zülpicher Strasse 49a, D-50674 Köln, Germany; preussef@uni-koeln.de; HU.Kasper@uni-koeln.de ²Geologisches Institut, Universität Bern, Baltzerstrasse 1, CH-3012 Bern, Switzerland.

(Received 5 February 2001; in final form 24 April 2001)

Introduction

Determination of dose rates for luminescence and electron spin resonance (ESR) dating is possible using a variety of different analytic methods. The investigations presented here were initiated by the need to determinate the concentration of dose rate relevant elements (K, Th, U) of sediment samples from drilling cores. High resolution (HR) gamma spectrometry, predominantly used in our laboratory, was not practical due to the limited amounts of sample material available. One major advantage of inductively coupled plasma - mass spectrometry (ICP-MS) is the possibility to analyse samples as small as 10 mg. The performance of sample preparation (homogeneity, digestion) and of the measurement itself can be verified by comparison with certified reference materials (CRM).

Disequilibria within the thorium and uranium decay series is a potential error source in luminescence and ESR dating. The most suitable approach to test for disequilibria is a combination of alpha spectrometry and low-level gamma spectrometry (Krbetschek et al., 1994; Olley et al., 1996). However, the low activities of ²³²Th and ²³⁸U make these measurements rather time consuming (some days per sample) precluding regular checks in routine dating procedures. An alternative method to allow evaluation of sample secular equilibrium is to measure the mother isotope at the head (ICP-MS) and the daughters at the end of a decay chain (HR gamma spectrometry). However, it is first necessary to exclude systematic differences between the two methods caused by difficulties with analytical procedures.

The results of ICP-MS and HR gamma spectrometry are compared for a variety of samples from different geological sites. These samples are

likely to be in equilibrium state due to their depositional history. For one sample, repeated measurements were carried out to obtain more information concerning the reproducibility of the results. A brief description of samples used in this study is presented in Tab. 1.

Table 1.

Description of samples analysed within this study. More information about the certified reference materials (CRM) is given by (*1) Govindaraju (1994) and (*2) Pszonicki et al. (2000), respectively

Sample	Origin	Sediment type	Grain-size	
AJS 1–3	Oman	aeolianite	sand	
BBR	Germany	house standard (basalt)		
BCR-1	USA	CRM (basalt) *1		
BGR 1-	Argentina	loess and dune sand	silt	
11			or sand	
GSD-3	China	CRM (fluvial) *1		
GSD-4	China	CRM (fluvial) *1		
HAB 1-3	Switzerland	fluvial deposit	sand	
			or silt	
HÜW 1-3	Switzerland	overbank deposit	sand	
			or silt	
NUSSY	Germany	loess	silt	
PGM 1-4	New	estuary deposit	sandy silt	
	Zealand			
PIC 5-7	New	coastal deposit	sand or	
	Zealand		silt	
SOIL-7	Austria	CRM (soil) *2		
WIL 1-5	Oman	desert lake and dune	sand	
		sand		

High resolution (HR) gamma spectrometry

HR gamma spectrometry was carried out using a high-purity germanium detector (Canberra GC2019–7935.2, coaxial p/n-type, 20 % relative efficiency, 1.9 keV FWHM at 1.33 MeV). After drying the

samples at 50°C and determining natural water content, 800 g or 1600 g of sediment were filled into Marinelli beakers. Analyses were performed after one month of storage to build up equilibrium between radon and its daughters. Samples were measured for 20 hours. The influence of environmental radioactivity was minimised by a shielding of low-activity lead. Using large samples and measuring for a long time significantly reduces the effect of background variability. Since natural radioactivity is generally weak for the samples which had been analysed $(0.5 - 2.0 \% \text{ of K}, 1 - 15 \mu \text{g g}^{-1} \text{ of }$ Th, $1-4 \ \mu g \ g^{-1}$ of U), most of the thorium and uranium series y-lines are close to the detection limit. Calibration was done by home standards containing known quantities of added radioactive elements (IAEA).

The results shown in Tab. 2 represent the average of 11 measurements of sample NUSSY, an Upper Weichselian loess from the Nußloch quarry near Heidelberg. All analyses were carried out during a period of two years using the same sealed container. This procedure provides accurate information of uncertainty due to peak matching and the error caused by background variability. The relative standard deviation (RSD) for different γ -lines ranges from approximately 1 % to 6 %. The poorer reproducibility is given for energy peaks where the signal/background ratio is low (e.g. for the peak of ²¹²Pb at 300.1 keV). The RSD of the average of all measurements and of all energy peaks is 2.1 % for K, 4.6 % for Th and 3.4 % for U.

Inductively coupled plasma – mass spectrometry (ICP-MS)

50 g of each sample were pulverised in a laboratory disc mill (tungsten carbide grinding barrel) and homogenised for analysis. Prior to ICP--MS analysis the sample material was dried overnight at 105°C (dry basis). All reagents used were Suprapur[®] grade (Merck).

Using conventional HF/HNO₃ decomposition, the recovery of Rb, Th and U will be severely restrained by selective coprecipitation with insoluble fluorides of major elements. The use of $HClO_4$ instead of HNO_3 enables an almost 100 % recovery of trace elements by completely suppressing fluoride precipitation (Yokoyama et al., 1999). Quantitative decomposition of samples, especially those containing resistant heavy minerals (e.g. Zircon), was considered using two different digestion protocols:

(1) Conventional open system decomposition

- 3 ml of 10 M HClO₄ and 3 ml of 22 M HF were added slowly to the weighed sample (15 ml Savilex PFA-vials, 100 mg) to prevent losses by violent carbonate reaction. After repeated sequential evaporation with 6 M HCl and 5.5 M HNO₃ the residue was dissolved with 0.5 M HNO₃, transferred to 50 ml volumetric flasks and stored in tightly closed HD-PE bottles. This procedure lasts for at least 2–3 days.
- (2) High pressure MULTIWAVE microwave digestion system

This automated system is equipped with special designed 50 ml TFM-vessels in combination with an evaporation rotor (Anton-Paar/Perkin-Elmer 6EVAP) and an aspiration-evaporation device (Prolabo ASPIVAP) (Knapp et al., 1997). The p-T monitored microwave-powered closed system reaches maximum temperatures of 260° C (for 20 minutes) and a pressure of 70 bar. Reagents and sample quantity used are identical to the ones used in the conventional decomposition protocol. The main advantage of microwave digestion is the short time needed (ca. 7-8 hours) and a guaranteed quantitative sample dissolution (Kasper and Preusser, 2000).

The elements K, Rb, Th and U were analysed by inductively coupled plasma – quadrupole mass spectrometry (ICP–QMS, Sciex/Perkin-Elmer ELAN 6000). Special attention was focused on automatic and customised corrections for isotopic interference, interfering molecular species, specially customised resolution for K and internal standardisation.

The NUSSY sample was used for an interlaboratory comparison and the results are shown in Tab. 3. For most of the measurements, the results are similar when the standard deviation is considered. However, the contents of U analysed with ICP-MS at the Cologne laboratory are slightly lower compared with the average results determined by the other laboratories. The concentrations of K and Th analysed by a commercial laboratory are slightly higher in comparison with the contents measured by the other laboratories.

Comparison of ICP-MS, HR-gamma spectrometry and CRM

Concentrations of dose rate relevant elements of 34 samples have been determined using ICP-MS and HR gamma spectrometry, respectively (Fig. 1-3).





Comparison of concentration of K determined by HR gamma spectrometry (closed circles) and CRM (open circles), respectively, versus ICP-MS.





Comparison of concentration of Th determined by HR gamma spectrometry (closed circles) and CRM (open circles), respectively, versus ICP-MS.





Comparison of concentration of U determined by HR gamma spectrometry (closed circles) and CRM values (open circles), respectively, versus ICP-MS.





Plot of dose rates calculated for HR gamma spectrometry (closed circles) and CRM values (open circles), respectively, versus dose rates determined using the ICP-MS results.

Table 2.

HR gamma spectrometry results for the different energy peaks of potassium and of isotopes from the thorium and uranium series of the NUSSY sample. The averages (n = 11) of the deduced contents of dose rate relevant elements, standard deviation (SD) and relative standard deviation (RSD), and the signal to background (BG) ratio with standard deviation are shown for repeated measurements.

Element	Peak (keV)	Content	SD	RSD	signal/BG
Potassium (%)					
K-40	1460.0	0.96	0.02	2.1	26.3 ± 1.9
Thorium ($\mu g g^{-1}$)					
T1-208	583.2	7.29	0.18	2.5	3.5 ± 0.2
Tl -208	2614.5	7.25	0.27	3.7	40.2 ± 7.3
Pb-212	238.6	7.37	0.08	1.1	3.8 ± 0.1
Pb-212	300.1	7.43	0.47	6.3	0.37 ± 0.03
Bi-212	727.3	7.30	0.44	6.0	1.0 ± 0.1
Ac-228	338.3	7.67	0.35	4.6	1.3 ± 0.2
Ac-228	911.1	7.58	0.25	3.3	4.2 ± 0.2
Mean	_	7.41	0.34	4.6	-
Uranium (µg g ⁻¹)					
Pb-214	295.2	2.67	0.06	2.2	2.3 ± 0.1
Pb-214	351.9	2.70	0.04	1.5	4.6 ± 0.4
Bi-214	609.3	2.70	0.03	1.1	6.3 ± 0.6
Bi-214	1120.3	2.69	0.09	3.3	2.0 ± 0.2
Ra-226, U-235	186.0	2.61	0.16	6.1	0.53 ± 0.04
Mean	_	2.68	0.09	3.4	

Table 3.

Comparison of analyses determining the dose relevant elements (RSD) for the NUSSY sample carried out by different laboratories using ICP-MS and HR gamma spectrometry, respectively.

		K	Rb	Th	U
Laboratory		(%)	(µg g ⁻¹)	$(\mu g g^{-1})$	$(\mu g g^{-1})$
Cologne (y-spec.)	(n = 11)	0.96 (2.1)		7.4 (4.6)	2.7 (3.4)
Cologne (ICP-MS)	(n = 10)	1.04 (8.9)	51.8 (2.3)	7.7 (2.1)	2.3 (4.4)
GFZ (ICP-MS)	(n = 3)	_	51.6 (1.5)	7.7 (5.3)	2.6 (0.8)
Commercial (ICP-MS)	(n =1)	1.14	_	8.8	2.7

Generally, the results for both methods are in good agreement.

For some samples significantly higher K contents were determined using ICP-MS (Fig. 1). This is due to the atomic weight of K and hence the massive isotopic interference of Argon from the plasma source. With respect to reproducibility and accuracy of the ICP-MS results, we intend to carry out further tests using alternative analytical methods such as atomic absorption spectrometry (AAS) or flame emission spectrometry (FES). Additionally, the remarkable covariance of Rb and K might be used for cross-checking the results.

Slightly lower U contents in comparison with HR gamma spectrometry were measured using ICP-MS (Fig. 3) as it has already been shown in the laboratory inter-comparison test (Tab. 3). It is possible that the

conventional digestion protocol might not always guarantee complete dissolution of samples due to chemically resistant minerals such as Zircon. Thus, we favour the use of high-pressure microwave digestion in future research.

A comparison of dose rates calculated from the concentration of radioactive elements as determined by the two different methods is shown in Fig. 4. Dose rates have been calculated for $150 \,\mu\text{m}$ quartz grains assuming an alpha efficiency of 0.1 and a sediment moisture of 20 %. Neither cosmic rays nor internal dose rate contribution have been considered.

Calculations were carried out using the AGE program developed by R. Grün which uses the conversion factors of Nambi and Aitken (1986). The differences in dose rates of the two different methods is generally less than 10 % but it has to be considered that some of the ICP-MS results may be affected by methodological difficulties.

Conclusions

Determination of the concentration of dose rate relevant elements analysed by ICP-MS and HR gamma spectrometry yield consistent results for most of the samples. The ICP-MS results are in good agreement with values of analysed CRM. However, digestion of samples and the custom resolution technique used for the determination of K have to be considered carefully. Methodological improvements may further increase the accuracy of ICP-MS analyses, in particular for K. It is emphasised that a combination of ICP-MS and HR gamma spectrometry may reduce the uncertainties in dose rate determination by the ability to routinely check for secular disequilibria within the thorium and uranium decay series.

Acknowledgements

The NUSSY sample was homogenised and provided by J. Heinicke, A. Lang, U. Rieser and L. Zöller. P. Dulski, GeoForschungs Zentrum (GFZ) Potsdam, contributed with three ICP–MS analyses of the NUSSY sample. M. Bateman provided results determined by a commercial ICP–MS laboratory. We want to express our special thanks to N. Willems who initially calibrated the HR gamma spectrometer used in this study and to M. Kelly for improving the English. M. Krbetschek is greatly thanked for reviewing the manuscript which benefited from his constructive comments.

References

- Govindaraju, K. (1994). 1994 compilation of working values and sample description for 383 geostandards. Geostandards Newsletter, 18: 1– 158.
- Kasper, H.U. and Preusser, F. (2000). Determination of K, Rb, Th and U by ICP-QMS for trapped charge dating of Quaternary sediments. Geoanalysis 2000, 4th International Conference on the Analysis of Geological and Environmental Materials: p. 65; Pont a Mousson, France.
- Knapp, G., Maichin, B. and Baumgartner, U. (1997).
 Störungen der ICP-Emissionsspektralanalyse durch organische Aufschlussrückstände. In: Vogt, C., Wennrich, R. and Werner, G. (Eds.): Colloquium Analytische Atomsprektroskopie (CANAS '97), 105–109.
- Krbetschek, M.R., Rieser, U., Zöller, L. and Heinicke, J. (1994). Radioactive disequilibria in palaedosimetric dating of sediments. *Radiation Measurements*, 24: 485–489.
- Nambi, K.S.V. and Aitken, M.J. (1986). Annual dose conversion factors for TL and ESR dating. Archeometrie, 28: 202–205.
- Olley, J.M., Murray, A. and Roberts, R.G. (1996). The effects of disequilibria in the Uranium and Thorium decay chains on burial dose rates in fluvial sediments. *Quaternary Science Reviews* (*Quaternary Geochronology*), 15: 751–760.
- Pszonicki, L., Hanna, A.N. and Suschny, O. (2000). Reference material IAEA-SOIL7. International Atomic Energy Agency, Analytic Control Services, 4 pp.
- Yokoyama, T., Makishima, A. and Nakamura, E. (1999). Evaluation of coprecipitation of incompatible trace elements with fluoride during silicate rock dissolution by acid digestion. *Chemical Geology*, 157: 175–187.

Reviewer

M. Krbetschek

Thesis abstract

Thesis title: Studies of quartz luminescence sensitivity relevant to dating and dosimetry Author: Chen Geng Date: May 2000 Supervisor: Dr. S.H. Li ; Examiners : Prof. A.G. Wintle and Dr. J.K.C. leung University: The university of Hong Kong

Quartz luminescence sensitivities, both OSL and 110°C TL, to irradiation dose were studied. The sensitivity is expressed as the ratio of luminescence centres (L) to non luminescence centres (R), and is explained using a model in respect of its changes with dose, heating and UV exposure. A modified model is introduced involving multiple R hole centres. It is found that only one type of R centre is involved in the OSL sensitivity change, and it is shared with the 110°C TL. However, more hole centres are involved in production of the 110°C TL.

Experiments involving annealing at various temperatures indicate that sensitivity changes of both OSL and 110°C TL signals are temperature dependent. At temperatures below 500°C, different filling and emptying of R and L centres is the dominant process and governs the sensitivity changes. There is a linear relationship between the OSL sensitivity and 110°C TL sensitivity with regeneration dose. Most holes can be transferred from R centres to L centres by annealing at 500°C for 10 minutes. Above 500°C, the sensitivity change results mainly from the destruction of R centres. The initial destruction of R centres happens at the first phase change of quartz at 573°C. At the second phase change of quartz at 870°C, the destruction of L centres occurs.

Isothermal decay experiments and quick heat experiments are applied to obtain the parameters of the OSL trap and the R centres. The lifetime of the OSL trap is found to be in a range from 10^8 to 10^9 years at 20°C. It is suggested that the OSL signal is stable enough for the dating of one million-year old samples. The OSL 325°C trap contributes an overwhelming majority of he OSL signal used in the OSL dating process. The lifetime of the R center is found to lie in the range from 1 to 30 ka at 20°C. The decay of R centres results in OSL sensitivity changes. The effect of sensitivity change must be considered in dating and dosimetry measurements.

Competition for electron between the OSL trap (325°C TL trap) and other traps is observed. The

electron traps shallower than OSL trap compete with the OSL trap, but the traps deeper than the OSL trap do not.

Thesis abstract

Thesis title: Development and application of luminescence dating to quaternary sediments from China Author: Jiafu Zhang Date: August 2000 Supervisor: Dr. S.H. Li and Dr. M-Y.W. Tso ; examiners : Prof. T.M. Chen and Dr. D.D. Zhang University: The university of Hong Kong

A three-stage single-aliquot additive-dose protocol (3-SAAD) equivalent dose for the (ED) determination of potassium feldspar is proposed. This procedure consists of 3 stages: preheat/measurement, dose/preheat/measurement and preheat/measurement; all the measurements for the equivalent dose determination are made on a single aliquot of feldspar. 3-SAAD provides a simple and efficient way of checking the appropriateness of preheat conditions during the ED estimation; the effects of the differences in mineralogy and luminescence properties from aliquot to aliquot are minimized. This procedure is also suitable for ED measurements on individual grains of feldspar.

An isochron technique is proposed, based on the EDs of grains (or aliquots) as a function of internal dose rate. This technique uses both quartz and feldspars grains (of the same size range), and takes advantage of the range in internal dose rates in these minerals to overcome uncertainties in the external dose rate. Even in the absence of independent age controls, the validity of the isochron ages can be assessed using the isochron equation. A significant improvement in dating uncertainty is demonstrated.

Assessment of the adequacy of bleaching of optical signals is very important for optical dating of sediments. A new method of assessing the bleaching of sedimentary feldspar grains is proposed, based on the difference in the bleaching rate between IRSL and TL signals. The size and scatter of the bleaching factors (normalized ratio of IRSL to TL) of grains (or aliquots) from a sample indicate the degree of bleaching. Well bleached grains can be identified by a relatively small bleaching factor, and relatively small scatter. All measurements required to estimate the ED and the degree of bleaching are carried out on the same grain (or aliquot). HF etching is necessary in optical dating of feldspar. Feldspars from different samples have different etching characteristics. For sedimentary feldspars from Hong Kong, it is found that 40 min etching with 10% HF at room temperature is appropriate.

Finally, three types of sediment from different areas are systemically dated using the optical techniques. They are colluvium sediments from Sham Wat debris lobe, Hong Kong, coastal deposits from area offshore of Hong Kong and dune sand samples from Horqin sandyland of northeastern China. Potential difficulties in optical dating of such sediments are discussed.

Note

Some notes on language

The language of science is supposedly precise. In practice it is often not so, sometimes through ignorance or carelessness, and sometimes because precision requires so much to be said as to be impracticable. With this in mind, I hope these notes will help curtail some of the misuses of terms that have crept into the literature. I write these notes knowing full well that I am sometimes guilty of comparable offences against the language.

Age or Date? A date refers to a specific year in the past, for example AD1666 or 326 BC. An age refers to the amount of time of past existence, for example an age of 350 years means the thing referred to has existed for 350 years. The word "date" is often used erroneously when "age" is meant. "Date lists" are often really "age lists".

Black-body radiation. This is an idealized concept. Imagine a closed cavity in thermodynamic equilibrium, then the electromagnetic radiation within the cavity is a function only of the temperature, and is referred to as black-body radiation. If a tiny hole is made in the cavity then the radiation that comes out of the hole is a close approximation to black-body radiation. In contrast the radiation that is emitted by something that is hot is NOT blackbody radiation unless it is perfectly "black" and no such material exists. Such radiation is of less intensity than black-body radiation, and the spectrum will be different. The word incandescence is used to describe the emission from a tungsten light bulb or any other visibly hot object.

Dose.

The term "radiation dose" has a well-defined meaning; it is the amount of radiation energy absorbed per unit mass of matter. Thus if 1 joule of radiation energy is absorbed in 1 kilogram of matter, the radiation dose is 1 $J.kg^{-1}$ or 1 Gy. The key words here are 'energy' and 'absorbed'. What happens to the energy after it is absorbed is completely irrelevant to the evaluation of the dose. In practice, most of the energy will end up in the form of heat and leave the matter in question by thermal conduction, convection or electromagnetic radiation (there is an exception to this is if the matter is thermally isolated from the surroundings). A small fraction of the energy absorbed will end up stored by electrons and atoms in excited states. It would be wrong to talk about the absorbed radiation energy contained in something. Thus it would be equally wrong to talk about a dose contained in something and the phrases "dose in", "stored dose", "contained dose", acquired dose" and such should not be used.

Dose rate or Annual Dose? The annual dose is the radiation dose that occurs in one year and the SI unit for it is the gray (Gy). The dose rate is the radiation dose per unit time, and the practical unit for it is Gy.ka⁻¹. The term "annual dose rate" should not be used.

Luminescence. Luminescence is the light emitted by a substance in response to a stimulus. The stimulus may be heat, light, sound, shock, a beam of electrons, or anything else. The terms thermoluminescence, photoluminescence, sono-luminescence etc are used to refer to the light emitted in the cases of different stimuli. The term "optical luminescence" that is sometimes used does not make any sense since luminescence is already an optical phenomenon. Luminescence is not something that can be stored, removed, reset or set to zero.

Palaeodose. This word refers to a past radiation dose. The difficulty I have with current use is that in our work it is not the actual past radiation dose that is determined, but the beta or gamma dose that results in the same luminescence intensity during thermal or optical excitation. In the past, the term equivalent dose was introduced to deal with this difference. The introduction of the term was necessary because there can be a large difference between the actual radiation dose and the equivalent dose. This difference arises mainly because alpha particles produce different effects gammas do betas than or on а per unit-of-absorbed-energy basis. Someone reading our literature could easily be misled into thinking that palaeodose meant the actual radiation dose. For this reason I think we should be using the term palaeodose equivalent, Pe or Peq. For the same reason we should not be using the term 'dose rate' but be using 'equivalent dose rate', since what we evaluate is not the true dose rate.

Hz and Bq. Herz (Hz) is the unit for the frequency of a periodic signal. This is one that repeats on a regular basis, such as the sound of a tuning fork or the electrical voltage at a wall socket in a house. 1 Hz is 1 cycle per second. Becquerel (Bq) is the SI unit of activity of a radioactive source. The decays occur randomly, not periodically, in time. An activity of 1 Bq means the decays occur at an average rate of 1 per second. It is not appropriate to use either of these for the rate at which photons are emitted from a sample, or the rate at which photons are counted. Photon emission is a random process, not periodic, thus precluding the use of Hz. It is not radioactive decay, thus precluding the use of Bq. Hence one should use counts per second or something similar as the unit for the photon count rate.

D.J.Huntley Physics Dept., Simon Fraser University, Burnaby, B. C., V5A 1S6, Canada

Bibliography

(From 1st October 2000 to 30th March 2001) Compiled by Ann Wintle

Akridge J. M. C., Benoit P. H., and Sears D. W. G. (2001) Determination of trapping parameters of the high temperature thermoluminescence peak in equilibrated ordinary chondrites. *Radiation Measurements* 33, 109-117.

Arkhipov S. A., Gnibidenko Z. N., and Shelkoplyas V. N. (2000) Correlation and paleomagnetism of glacial and loess-paleosol sequences on the West Siberian Plain. *Quaternary International* **68**, 13-27.

Bailey R. M. (2001) Towards a general kinetic model for optically and thermally stimulated luminescence of quartz. *Radiation Measurements* 33, 17-45.

Banerjee D. (2001) Supralinearity and sensitivity changes in optically stimulated luminescence of annealed quartz. *Radiation Measurements* 33, 47-57.

Banerjee D., Murray A. S., Bøtter-Jensen L., and Lang A. (2001) Equivalent dose estimation using a single aliquot of polymineral fine grains. *Radiation Measurements* 33, 73-94.

Berger G. W. and Péwé T. L. (2001) Last Interglacial age of the Eva Forest Bed, central Alaska, from thermoluminescence dating of bracketing loess. *Quaternary Science Reviews* 20, 484-498.

Blackwell B. A. B. and Blickstein J. I. B. (2000) Considering sedimentary uptake in external dose rate determinations for ESR and luminescent dating. *Quaternary International* 68, 329-343.

Blackwell B. A. B., Leung H. Y., Skinner A. R., Schwarcz H. P., Lebel S., Valladas H., Blickstein J. I. B., and Divjak M. N. (2000) External dose rate determinations for ESR dating at Bau de l'Aubesier, Provence, France. *Quaternary International* 68, 345-361.

Carter J. A. and Lian O. B. (2000) Palaeoenvironmental reconstruction from last interglacial using phytolith analysis, southeastern North Island, New Zealand. *Journal of Quaternary Science* 15, 733-743.

Chen G., Murray A. S., and Li S. H. (2001) Effect of heating on the quartz dose-response curve. Radiation Measurements 33, 59-63.

Chen R. and Leung P. L. (2000) A model for dose-rate dependence of thermoluminescence intensity. Journal of Physics D - Applied Physics 33, 846-850.

Chruscinska A. (2001) The fractional thermoluminescence of bleached K-feldspars. Journal of Luminescence 92, 271-276.

Clemmensen L. B., Pye K., Murray A., and Heinemeier J. (2001) Sedimentology, stratigraphy and landscape evolution of a Holocene coastal dune system, Lodbjerg, NW Jutland, Denmark. Sedimentology 48, 3-27.

Collen J. D., Grapes R. H., and Eagar S. H. (2000) Moa fossils and chronology of a Pleistocene terrace sequence, Tauweru, Wairarapa, New Zealand. New Zealand Journal of Geology and Geophysics 43, 483-491.

Eriksson M. G., Olley J. M., and Payton R. W. (2000) Soil erosion history in central Tanzania based on OSL dating of colluvial and alluvial hillslope deposits. *Geomorphology* 36, 107-128.

Galloway R. B. (2001) Luminescence dating: limitations to accuracy attainable. Journal of Radioanalytical and Nuclear Chemistry 247, 679-683.

Goudie A. S., Colls A., Stokes S., Parker A., White K., and Al-Farraj A. (2000) Latest Pleistocene and Holocene dune construction at the north-eastern edge of the Rub Al Khali, United Arab Emirates. *Sedimentology* 47, 1011-1021.

Grün R., Moriarty K., and Wells R. (2001) Electron spin resonance dating of the fossil deposits in the Naracoorte Caves, South Australia. *Journal of Quaternary Science* 16, 49-59.

Grün R. and Stringer C. (2000) Tabun revisited: revised ESR chronology and new ESR and U-series analyses of dental material from Tabun C1. Journal of Human Evolution 39, 601-612.

Hashimoto T., Komatsu Y., Hong D. G., and Uezu Y. (2001) Determination of radionuclides in small pieces of archaeological sample and its application to TL-dating. *Radiation Measurements* 33, 95-101.

Hatté C., Antoine P., Fontugne M., Lang A., Rousseau D.-D., and Zöller L. (2001) Delta 13C of loess organic matter as a potential proxy for paleoprecipitation. *Quaternary Research* 55, 33-38.

Helmens K. F., Rasanen M. E., Johansson P. W., Jungner H., and Korjonen K. (2000) The Last Interglacial-Glacial cycle in NE Fennoscandia: a nearly continuous record from Sokli (Finnish Lapland). *Quaternary Science Reviews* **19**, 1605-1623.

Hong D. G., Galloway R. B., and Hashimoto T. (2001) Equivalent dose determination from quartz additive dose optically stimulated luminescence: simulation tests. *Nuclear Instruments and Methods in Physics Research B* 173, 533-540.

Hong D.-G., Yi S. B., Galloway R. B., Tsuboi T., and Hashimoto T. (2001) Optical dating of archaeological samples using a single aliquot of quartz stimulated by blue light. *Journal of Radioanalytical and Nuclear Chemistry* 247, 179-184.

Hütt G., I. J., and Vasilchenko V. (2001) Photoionization of radiation-induced traps in quartz and alkali feldspars. Applied Radiation and Isotopes 54, 175-182.

Kar A., Singhvi A. K., Rajaguru S. N., Juyal N., Thomas J. V., Banerjee D., and Dhir R. P. (2001) Reconstruction of the late Quaternary environment of the lower Luni plains, Thar Desert, India. *Journal of Quaternary Science* 16, 61-68.

Kaufman D. S., Manley W. F., Forman S. L., and Layer P. W. (2001) Pre-Late-Wisconsin glacial history, coastal Ahklun Mountains, southwestern Alaska - new amino acid, thermoluminescence, and 40Ar/39Ar results. *Quaternary Science Reviews* 20, 337-352.

Lehmkuhl F., Klinge M., Rees-Jones J., and Rhodes E. J. (2000) Late Quaternary aeolian sedimentation in central and south-eastern Tibet. *Quaternary International* 68, 117-132.

Lehmkuhl F. and Lang A. (2001) Geomorphological investigations and luminescence dating in the southern part of the Khangay and the Valley of the Gobi Lakes (Central Mongolia). *Journal of Quaternary Science* 16, 69-87.

Lian O., Wang N. S., Grapes R. H., and Collen J. D. (2000) Optical dating of sediments associated with a fossil moa skeleton near Tauweru, Wairarapa, New Zealand. New Zealand Journal of Geology and Geophysics 43, 493-499.

Lian O. B. and Shane P. A. (2000) Optical dating of paleosols bracketing the widespread Rotoehu tephra, North Island, New Zealand. *Quaternary Science Reviews* 19, 1649-1662.

Mahaney W. C., Milner M. W., Voros J., Kalm V., Hutt G., Bezada M., Hancock R. G. V., and Aufreiter S. (2000) Stratotype for the Merida Glaciation at Pueblo Llano in the northern Venezuelan Andes. *Journal of South American Earth sciences* 13, 761-774.

Mauz B. and Hassler U. (2000) Luminescence chronology of Late Pleistocene raised beaches in southern Italy: new data of relative sea-level changes. *Marine Geology* **170**, 187-203.

Ohta M., Yasuda M., and Takami M. (2000) Thermoluminescence properties of SiO2:Al Eu. Journal of Alloys and Compounds 303, 320-324.

Oke G. and Yurdatapan E. (2000) Optically stimulated luminescence dating of pottery from Turkey. Talanta 53, 115-119.

Parkash B., Kumar S., Rao M. S., Giri S. C., Kumar C. S., Gupta S., and Srivastava P. (2000) Holocene tectonic movements and stress field in the western Gangetic plains. *Current Science* **79**, 438-449.

Rees-Jones J., Rink W. J., Norris R. J., and Litchfield N. J. (2000) Optical luminescence dating of uplifted marine terraces along the Akatore Fault near Dunedin, South Island, New Zealand. New Zealand Journal of Geology and Geophysics 43, 419-424.

Richards B. W. M., Benn D. I., Owen L. A., Rhodes E. J., and Spencer J. Q. (2000) Timing of late Quaternary glaciations south of Mount Everest in the Khumbu Himal, Nepal. *Geological Society of American Bulletin* **112**, 1621-1632.

Schellmann G. and Radtke U. (2000) ESR dating stratigraphically well-constrained marine terraces along the Patagonian Atlantic coast (Argentina). Quaternary International 68, 261-273.

Shinno I., Nakamura T., and Sekine T. (2000) Deformation-sensitive luminescence of forsterite (Mg-2 SiO4) shocked up to 82 CPa. *Journal of Luminescence* 87-9, 1292-1294.

Soika C. and Delincee H. (2000) Thermoluminescence analysis for detection of irradiated food - effects of dose rate on the glow curves of quartz. Lebensmittel-Wissenschaft und-Technologie 33, 440-443.

Soika C. and Delincee H. (2000) Thermoluminescence analysis for detection of irradiated food - luminescence characteristics of minerals for different types of radiation and radiation doses. *Lebensmittel-Wissenchaft und-Technologie* 33, 431-439.

Srivastava P., Juyal N., Singhvi A. K., Wasson R. J., and Bateman M. D. (2001) Luminescence chronology of river adjustment and incision of Quaternary sediments in the alluvial plain of the Sabarmati River, north Gujurat, India. *Geomorphology* **36**, 217-229.

Trautmann T. (2000) A study of radioluminescence kinetics of natural feldspar dosimeters: experiments and simulations. Journal of Physics D - Applied Physics 33, 2304-2310.

Turney C. S. M., Bird M. I., Fifield L. K., Roberts R. G., Smith M., Dortch C. E., Grün R., Lawson E., Ayliffe L. K., Miller G. H., Dortch J., and Cresswell R. G. (2001) Early human occupation at Devil's Lair, southwestern Australia 50,000 years ago. *Quaternary Research* 55, 3-13.

Wilson P., Orford J. D., Knight J., Braley S. M., and Wintle A. G. (2001) Late-Holocene (post-45000 years BP) coastal dune development in Northumberland, northeast England. *The Holocene* 11, 215-229.

Wolfe A. P., Frechette B., Richard P. J. H., Miller G. H., and Forman S. L. (2000) Paleoecology of a > 90,000-year lacustrine sequence from Fog Lake, Baffin Island, Arctic Canada. *Quaternary Science Reviews* 19, 1677-1699.

Wolfe S. A., Huntley D. J., David P. P., Ollerhead J., Sauchyn D. J., and MacDonald G. M. (2001) Late 18th century drought-induced sand dune activity, Great Sand Hills, Saskatchewan. *Canadian Journal of Earth Sciences* **38**, 105-117.

Yin G. M. and Li S. H. (2000) Thermoluminescence dating of the last volcanic eruption in Changbai Mountain. Acta Petrologica Sinica 16, 353-356.

Zhang J. F., Li S. H., and Tso M. Y. W. (2001) Assessment of bleaching of K feldspar-grains. Radiation Measurements 33, 103-108.

Zhang J. F., Li S. H., and Tso M. Y. W. (2001) Improvement of the equivalent dose determination using aliquots of potassium feldspar. *Radiation Measurements* 33, 65-71.

Zhou L. P., van Andel T. H., and Lang A. (2000) A luminescence dating study of open-air palaeolithic sites in western Epirus, Greece. Journal of Archaeological Science 27, 609-620.

Zilberman E., Amit R., Heimann A., and Porat N. (2000) Changes in Holocene Paleoseismic activity in the Hula pull-apart basin, Dead Sea Rift, northern Israel. *Tectonophysics* 321, 237-252.

Papers from Rome LED99 volume of Quaternary Science Reviews in alphabetical order

Anthony I. M. C., Sanderson D. C. W., Cook G. T., Abernethy D., and Housley R. A. (2001) Dating a burnt mound from Kilmartin, Argyll, Scotland. *Quaternary Science Reviews* 20(5-9), 921-925.

Bailey S. D., Wintle A. G., Duller G. A. T., and Bristow C. S. (2001) Sand deposition during the last millennium at Aberffraw, Anglesey, North Wales as determined by OSL dating of quartz. *Quaternary Science Reviews* 20(5-9), 701-704.

Balescu S., Lamothe M., Auclair M., and Shilts W. W. (2001) IRSL dating of Middle Pleistocene interglacial sediments from southern Quebec (Canada) using multiple and single grain aliquots. *Quaternary Science Reviews* **20**(5-9), 821-824.

Banerjee D., Murray A. S., and Foster I. D. L. (2001) Scilly Isles, UK: optical dating of a possible tsunami deposit from the 1755 Lisbon earthquake. *Quaternary Science Reviews* **20**(5-9), 715-718.

Bateman M. D. and Herrero A. D. (2001) The timing and relation of aeolian sand deposition in central Spain to the aeolian sand record of NW Europe. *Quaternary Science Reviews* 20(5-9), 779-782.

Bitinas A., Damusyte A., Hutt G., Jaek I., and Kabailiene M. (2001) Application of the OSL dating for stratigraphic correlation of Late Weichselian and Holocene sediments in the Lithuanian Maritime Region. *Quaternary Science Reviews* 20(5-9), 767-772.

Blackwell B. A. B., Skinner A. R., and Blickstein J. I. B. (2001) ESR isochron exercises: how accurately do modern dose rate measurements reflect paleodose rates? *Quaternary Science Reviews* **20**(5-9), 1031-1039.

Bluszcz A. (2001) Simultaneous OSL and TL dating of sediments. Quaternary Science Reviews 20(5-9), 761-766.

Bonde A., Murray A., and Friedrich W. L. (2001) Santorini: Luminescence dating of a volcanic province using quartz? *Quaternary Science Reviews* 20(5-9), 789-793.

Chen Y. G., Wu W. S., Chen C. H., and Liu T. K. (2001) A date for volcanic eruption inferred from a siltstone xenolith. *Quaternary Science Reviews* 20(5-9), 869-873.

Chiavari C., Martini M., Sibilia E., and Vandini M. (2001) Thermoluminescence (TL) characterisation and dating feasibility of ancient glass mosaic. *Quaternary Science Reviews* 20(5-9), 967-972.

Clarke M. L., Rendell H. M., Hoare P. G., Godby S. P., and Stevenson C. R. (2001) The timing of coversand deposition in northwest Norfolk, UK: a cautionary tale. *Quaternary Science Reviews* 20(5-9), 705-713.

Colls A. E., Stokes S., Blum M. D., and Straffin E. (2001) Age limits on the late quaternary evolution of the upper Loire River. *Quaternary Science Reviews* 20(5-9), 743-750.

Engelmann A., Neber A., Frechen M., Boenigk W., and Ronen A. (2001) Luminescence chronology of Upper Pleistocene and Holocene aeolianites from Netanya South - Sharon Coastal Plain, Israel. *Quaternary Science Reviews* 20(5-9), 799-804.

Feathers J. K. and Migliorini E. (2001) Luminescence dating at Katanda - a reassessment. Quaternary Science Reviews 20(5-9), 961-966.

Fehrentz M. and Radtke U. (2001) Luminescence dating of Pleistocene outwash sediments of the Senne area (Eastern Munsterland, Germany). Quaternary Science Reviews 20(5-9), 725-729.

Folz E., Bodu P., Bonte P., Joron J. L., Mercier N., and Reyss J. L. (2001) OSL dating of fluvial quartz from Le Closeau, a Late Paleolithic site near Paris - comparison with C-14 chronology. *Quaternary Science Reviews* 20(5-9), 927-933.

Frechen M., Dermann B., Boenigk W., and Ronen A. (2001) Luminescence chronology of aeolianites from the section at Givat Olga - Coastal Plain of Israel. *Quaternary Science Reviews* 20(5-9), 805-809.

Fuchs M. and Lang A. (2001) OSL dating of coarse-grain fluvial quartz using single-aliquot protocols on sediments from NE Peloponnese, Greece. *Quaternary Science Reviews* **20**(5-9), 783-787.

Fukuchi T. (2001) Assessment of fault activity by ESR dating of fault gouge; an example of the 500 m core samples drilled into the Nojima earthquake fault in Japan. *Quaternary Science Reviews* **20**(5-9), 1005-1008.

Gautier A. (2001) Luminescence dating of archaeometallurgical slag: use of the SAR technique for determination of the burial dose. *Quaternary Science Reviews* **20**(5-9), 973-980.

Hagihara N., Miono S., Manabe S., and Nakanishi A. (2001) A discovery of suitable layers for the thermoluminescence (TL) dating of ancient iron manufacturing furnace to investigate the origin of iron manufacturing in Japan. *Quaternary Science Reviews* **20**(5-9), 987-991.

Han Z. Y., Li S. H., and Tso M. Y. W. (2001) TL dating of granitic quartz using an additive alpha dose method. *Quaternary Science Reviews* 20(5-9), 907-911.

Haustein M., Krbetschek M. R., Trautmann T., Roewer G., and Stolz W. (2001) A luminescence study for dating archaeometallurgical slags. *Quaternary Science Reviews* 20(5-9), 981-985.

Hilgers A., Murray A. S., Schlaak N., and Radtke U. (2001) Comparison of quartz OSL protocols using lateglacial and Holocene dune sands from Brandenburg, Germany. *Quaternary Science Reviews* **20**(5-9), 731-736.

Hoffmann D., Woda C., Strobl C., and Mangini A. (2001) ESR-dating of the Arctic sediment core PS1535 doseresponse and thermal behaviour of the CO2--signal in foraminifera. *Quaternary Science Reviews* 20(5-9), 1009-1014.

Huntley D. J. and Prescott J. R. (2001) Improved methodology and new thermoluminescence ages for the dune sequence in south-east South Australia. *Quaternary Science Reviews* **20**(5-9), 687-699.

Hutt G., Goksu H. Y., Jaek I., and Hiekkanen M. (2001) Luminescence dating of Somero sacristy, SW Finland using the 210 degrees C TL peak of quartz. *Quaternary Science Reviews* 20(5-9), 773-777.

Jungner H., Korjonen K., Heikkinen O., and Wiedemann A. M. (2001) Luminescence and radiocarbon dating of a dune series at Cape Kiwanda, Oregon, USA. *Quaternary Science Reviews* 20(5-9), 811-814.

Kuzmin Y. V., Hall S., Tite M. S., Bailey R., O'Malley J. M., and Medvedev V. E. (2001) Radiocarbon and thermoluminescence dating of the pottery from the early Neolithic site of Gasya (Russian Far East): initial results. *Quaternary Science Reviews* 20(5-9), 945-948.

Lang A. and Zolitschka B. (2001) Optical dating of annually laminated lake sediments - A test case from Holzmaar/Germany. *Quaternary Science Reviews* 20(5-9), 737-742.

Lee H. K. and Schwarcz H. P. (2001) ESR dating of the subsidiary faults in the Yangsan fault system, Korea. *Quaternary Science Reviews* 20(5-9), 999-1003.

Li S. H. and Yin G. M. (2001) Luminescence dating of young volcanic activity in China. Quaternary Science Reviews 20(5-9), 865-868.

Miallier D., Gibert F., Fain J., Pilleyre T., and Sanzelle S. (2001) Fluid inclusions in quartz: interference with thermoluminescence and its application to dating. *Quaternary Science Reviews* 20(5-9), 901-905.

Molodkov A. (2001) ESR dating evidence for early man at a Lower Palaeolithic cave- site in the Northern Caucasus as derived from terrestrial mollusc shells. *Quaternary Science Reviews* **20**(5-9), 1051-1055.

Murray A. S. and Clemmensen L. B. (2001) Luminescence dating of Holocene aeolian sand movement, Thy, Denmark. *Quaternary Science Reviews* 20(5-9), 751-754.

Ninagawa K., Kitahara T., Toyoda S., Hayashi K., Nishido H., Kinjo M., and Kawana T. (2001) Thermoluminescence dating of the Ryukyu Limestone. *Quaternary Science Reviews* 20(5-9), 829-833.

Pike A. W. G. and Hedges R. E. M. (2001) Sample geometry and U uptake in archaeological teeth: implications for U-series and ESR dating. *Quaternary Science Reviews* 20(5-9), 1021-1025.

Porat N., Zilberman E., Amit R., and Enzel Y. (2001) Residual ages of modern sediments in an hyperarid region, Israel. *Quaternary Science Reviews* 20(5-9), 795-798.

Price D. M., Brooke B. P., and Woodroffe C. D. (2001) Thermoluminescence dating of aeolianites from Lord Howe Island and South-West Western Australia. *Quaternary Science Reviews* 20(5-9), 841-846.

Radtke U., Janotta A., Hilgers A., and Murray A. S. (2001) The potential of OSL and TL for dating Late glacial and Holocene dune sands tested with independent age control of the Laacher See tephra (12 880 a) at the Section 'Mainz-Gonsenheim'. *Quaternary Science Reviews* 20(5-9), 719-724.

Rich J. and Stokes S. (2001) Optical dating of geoarchaeologically significant sites from the Southern High Plains and South Texas, USA. *Quaternary Science Reviews* **20**(5-9), 949-959.

Richardson C. A. (2001) Residual luminescence signals in modern coastal sediments. Quaternary Science Reviews 20(5-9), 887-892.

Rink W. J. and Pieper K. D. (2001) Quartz thermoluminescence in a storm deposit and a welded beach ridge. *Quaternary Science Reviews* 20(5-9), 815-820.

Roberts H. M. and Wintle A. G. (2001) Equivalent dose determinations for polymineralic fine-grains using the SAR protocol: application to a Holocene sequence of the Chinese Loess Plateau. *Quaternary Science Reviews* 20(5-9), 859-863.

Rogue C., Guibert P., Vartanian E., Bechtel F., and Schvoerer M. (2001) Thermoluminescence-dating of calcite: study of heated limestone fragments from Upper Paleolithic layers at Combe Sauniere, Dordogne, France. *Quaternary Science Reviews* 20(5-9), 935-938.

Sanderson D. C. W., Bishop P., Houston I., and Boonsener M. (2001) Luminescence characterisation of quartz-rich cover sands from NE Thailand. *Quaternary Science Reviews* **20**(5-9), 893-900.

Schellmann G. and Radtke U. (2001) Progress in ESR dating of Pleistocene corals - a new approach for D-E determination. *Quaternary Science Reviews* **20**(5-9), 1015-1020.

Schwarcz H. P. and Rink W. J. (2001) Skinflint dating. Quaternary Science Reviews 20(5-9), 1047-1050.

Shlukov A. I., Usova M. G., Voskovskaya L. T., and Shakovets S. A. (2001) New dating techniques for Quaternary sediments and their application on the Russian Plain. *Quaternary Science Reviews* 20(5-9), 875-878.

Skinner A. R., Chasteen N. D., Shao J. L., and Blackwell B. A. B. (2001) Q-band studies of the ESR signal in tooth enamel. *Quaternary Science Reviews* 20(5-9), 1027-1030.

Sommerville A. A., Sanderson D. C. W., Hansom J. D., and Housley R. A. (2001) Luminescence dating of aeolian sands from archaeological sites in Northern Britain: a preliminary study. *Quaternary Science Reviews* 20(5-9), 913-919.

Spooner N. A., Olley J. M., Questiaux D. G., and Chen X. Y. (2001) Optical dating of an aeolian deposit on the Murrumbidgee floodplain. *Quaternary Science Reviews* 20(5-9), 835-840.

Stokes S., Bray H. E., and Blum M. D. (2001) Optical resetting in large drainage basins: tests of zeroing assumptions using single-aliquot procedures. *Quaternary Science Reviews* 20(5-9), 879-885.

Strickertsson K., Murray A. S., and Lykke-Andersen H. (2001) Optically stimulated luminescence dates for Late Pleistocene Sediments from Stensnaes, Northern Jutland, Denmark. *Quaternary Science Reviews* 20(5-9), 755-759.

Tanaka K., Hataya R., Spooner N. A., and Questiaux D. G. (2001) Optical dating of river terrace sediments from Kanto plains, Japan. *Quaternary Science Reviews* **20**(5-9), 825-828.

Tiemei C., Qi C., Quan Y., and Hu Y. Q. (2001) The problems in ESR dating of tooth enamel of Early Pleistocene and the age of Longgupo hominid, Wushan, China. *Quaternary Science Reviews* 20(5-9), 1041-1045.

Toyoda S., Rink W. J., Yonezawa C., Matsue H., and Kagami T. (2001) In situ production of alpha particles and alpha recoil particles in quartz applied to ESR studies of oxygen vacancies. *Quaternary Science Reviews* 20(5-9), 1057-1061.

Valladas H., Mercier N., Froget L., Joron J. L., Reyss J. L., and Aubry T. (2001) TL dating of Upper Palaeolithic sites in the Coa Valley (Portugal). *Quaternary Science Reviews* 20(5-9), 939-943.

Watanuki T. and Tsukamoto S. (2001) A comparison of GLSL, IRSL and TL dating methods using loess deposits from Japan and China. *Quaternary Science Reviews* 20(5-9), 847-851.

Woda C., Mangini A., and Wagner G. A. (2001) ESR dating of xenolithic quartz in volcanic rocks. *Quaternary* Science Reviews 20(5-9), 993-998.

Zhou L. P. and Shackleton N. J. (2001) Photon-stimulated luminescence of quartz from loess and effects of sensitivity change on palaeodose determination. *Quaternary Science Reviews* 20(5-9), 853-857.

Papers from Rome LED99 volume of Radiation Measurements in alphabetical order

Adamiec G. (2000) Variations in luminescence properties of single quartz grains and their consequences for equivalent dose estimation. *Radiation Measurements* **32**(5-6), 427-432.

Akselrod M. S., Larsen N. A., and McKeever S. W. S. (2000) A procedure for the distinction between static and dynamic radiation exposures of personal radiation badges using pulsed optically stimulated luminescence. *Radiation Measurements* **32**(3), 215-225.

Armitage S. J., Duller G. A. T., and Wintle A. G. (2000) Quartz from southern Africa: sensitivity changes as a result of thermal pretreatment. *Radiation Measurements* **32**(5-6), 571-577.

Bailey R. M. (2000) Circumventing possible inaccuracies of the single aliquot regeneration method for the optical dating of quartz. *Radiation Measurements* **32**(5-6), 833-840.

Bailey R. M. (2000) The interpretation of quartz optically stimulated luminescence equivalent dose versus time plots. *Radiation Measurements* 32(2), 129-140.

Bailey R. M. (2000) The slow component of quartz optically stimulated luminescence. Radiation Measurements **32**(3), 233-246.

Bailey R. M., Adamiec G., and Rhodes E. J. (2000) OSL properties of NaCl relative to dating and dosimetry. *Radiation Measurements* **32**(5-6), 717-723.

Bailiff I. K. (2000) Characteristics of time-resolved luminescence in quartz. Radiation Measurements 32(5-6), 401-405.

Bailiff I. K., Bøtter-Jensen L., Correcher V., Delgado A., Goksu H. Y., Jungner H., and Petrov S. A. (2000) Absorbed dose evaluations in retrospective dosimetry: methodological developments using quartz. *Radiation Measurements* 32(5-6), 609-613.

Bailiff I. K. and Holland N. (2000) Dating bricks of the last two millennia from Newcastle upon Tyne: a preliminary study. *Radiation Measurements* **32**(5-6), 615-619.

Barnett S. M. (2000) Sampling pottery in luminescence dating studies. Radiation Measurements 32(5-6), 467-472.

Bortolot V. J. (2000) A new modular high capacity OSL reader system. Radiation Measurements 32(5-6), 751-757.

Bøtter-Jensen L., Bulur E., Duller G. A. T., and Murray A. S. (2000) Advances in luminescence instrument systems. *Radiation Measurements* **32**(5-6), 523-528.

Bøtter-Jensen L., Solongo S., Murray A. S., Banerjee D., and Jungner H. (2000) Using the OSL single-aliquot regenerative-dose protocol with quartz extracted from building materials in retrospective dosimetry. *Radiation Measurements* **32**(5-6), 841-845.

Brennan B. J. (2000) Systematic underestimation of the age of samples with saturating exponential behaviour and inhomogeneous dose distribution. *Radiation Measurements* **32**(5-6), 731-734.

Brennan B. J., Prestwich W. V., Rink W. J., Marsh R. E., and Schwarcz H. P. (2000) Alpha and beta dose gradients in tooth enamel. *Radiation Measurements* 32(5-6), 759-765.

Bulur E. (2000) A simple transformation for converting CW-OSL curves to LM-OSL curves. Radiation Measurements 32(2), 141-145.

Bulur E., Bøtter-Jensen L., and Murray A. S. (2000) Optically stimulated luminescence from quartz measured using the linear modulation technique. *Radiation Measurements* **32**(5-6), 407-411.

Chen G., Li S. H., and Murray A. S. (2000) Study of the 110 degrees C TL peak sensitivity in optical dating of quartz. *Radiation Measurements* **32**(5-6), 641-645.

Chen R., Leung P. L., and Stokes M. J. (2000) Apparent anomalous fading of thermoluminescence associated with competition with radiationless transitions. *Radiation Measurements* **32**(5-6), 505-511.

Chithambo M. L. and Galloway R. B. (2000) On luminescence lifetimes in quartz. Radiation Measurements 32(5-6), 621-626.

Chithambo M. L. and Galloway R. B. (2000) Temperature dependence of luminescence time-resolved spectra from quartz. *Radiation Measurements* 32(5-6), 627-632.

Correcher V., Garcia-Guinea J., and Delgado A. (2000) Influence of preheating treatment on the luminescence properties of adularia feldspar (KAISi3O8). *Radiation Measurements* **32**(5-6), 709-715.

Debuyst R., Frechen M., and Idrissi S. (2000) Problems encountered in a TL and ESR study of natural monohydrocalcite. *Radiation Measurements* **32**(5-6), 725-729.

Duller G. A. T., Bøtter-Jensen L., and Murray A. S. (2000) Optical dating of single sand-sized grains of quartz: sources of variability. *Radiation Measurements* **32**(5-6), 453-457.

Erfurt G., Krbetschek M. R., Trautmann T., and Stolz W. (2000) Radioluminescence (RL) behaviour of Al2O3 : C-potential for dosimetric applications. *Radiation Measurements* **32**(5-6), 735-739.

Fattahi M. and Stokes S. (2000) Extending the time range of luminescence dating using red TL (RTL) from volcanic quartz. *Radiation Measurements* **32**(5-6), 479-485.

Fattibene P., Aragno D., Onori S., and Pressello M. C. (2000) Thermal induced EPR signals in tooth enamel. *Radiation Measurements* **32**(5-6), 793-798.

Franklin A. D., Prescott J. R., and Robertson G. B. (2000) Comparison of blue and red TL from quartz. Radiation Measurements 32(5-6), 633-639.

Galloway R. B. (2000) The initial rate of loss of luminescence due to infrared stimulation of K-feldspars. *Radiation Measurements* **32**(5-6), 703-707.

Grun R. (2000) Methods of dose determination using ESR spectra of tooth enamel. *Radiation Measurements* **32**(5-6), 767-772.

Habermann J., Schilles T., Kalchgruber R., and Wagner G. A. (2000) Steps towards surface dating using luminescence. *Radiation Measurements* **32**(5-6), 847-851.

Han Z. Y., Li S. H., and Tso M. Y. W. (2000) Effects of annealing on TL sensitivity of granitic quartz. Radiation Measurements 32(3), 227-231.

Hayes R. B. and Haskell E. H. (2000) A method for identification of interfering signals in EPR dating of tooth enamel. *Radiation Measurements* **32**(5-6), 781-785.

Krbetschek M. R. and Trautmann T. (2000) A spectral radioluminescence study for dating and dosimetry. *Radiation Measurements* 32(5-6), 853-857.

Krbetschek M. R., Trautmann T., Dietrich A., and Stolz W. (2000) Radioluminescence dating of sediments: methodological aspects. *Radiation Measurements* **32**(5-6), 493-498.

Kuhn R., Trautmann T., Singhvi A. K., Krbetschek M. R., Wagner G. A., and Stolz W. (2000) A study of thermoluminescence emission spectra and optical stimulation spectra of quartz from different provenances. *Radiation Measurements* **32**(5-6), 653-657.

Kuhns C. K., Larsen N. A., and McKeever S. W. S. (2000) Characteristics of LM-OSL from several different types of quartz. *Radiation Measurements* 32(5-6), 413-418.

Lamothe M. and Auclair M. (2000) The fadia method: a new approach in luminescence dating using the analysis of single feldspar grains. *Radiation Measurements* **32**(5-6), 433-438.

Larsen N. A., Bulur E., Bøtter-Jensen L., and McKeever S. W. S. (2000) Use of the LM-OSL technique for the detection of partial bleaching in quartz. *Radiation Measurements* **32**(5-6), 419-425.

Lepper K., Larsen N. A., and McKeever S. W. S. (2000) Equivalent dose distribution analysis of Holocene eolian and fluvial quartz sands from Central Oklahoma. *Radiation Measurements* **32**(5-6), 603-608.

Malmberg R. and Radtke U. (2000) The alpha-efficiency of corals and its importance for ESR- dating. Radiation Measurements 32(5-6), 747-750.

Marsh R. E. and Rink W. J. (2000) Beta-gradient isochron dating of thin tooth enamel layers using ESR. Radiation Measurements 32(5-6), 567-570.

Martini M., Meinardi F., and Vedda A. (2000) The role of alkali ions in the 190 K TSL peak in quartz. Radiation Measurements 32(5-6), 673-677.

McCoy D. G., Prescott J. R., and Nation R. J. (2000) Some aspects of single-grain luminescence dating. *Radiation Measurements* 32(5-6), 859-864.

Murray A. S. and Wintle A. G. (2000) Application of the single-aliquot regenerative-dose protocol to the 375 degrees C quartz TL signal. *Radiation Measurements* **32**(5-6), 579-583.

Murray A. S. and Wintle A. G. (2000) Luminescence dating of quartz using an improved single-aliquot regenerative-dose protocol. *Radiation Measurements* 32(1), 57-73.

Niedermayer M., Goksu H. Y., Dalheimer A., and Bayer A. (2000) Infrared (IR) stimulated luminescence from modern bricks in retrospective dosimetry applications. *Radiation Measurements* **32**(5-6), 825-832.

Onori S., Aragno D., Fattibene P., Petetti E., and Pressello M. C. (2000) ISS protocol for EPR tooth dosimetry. *Radiation Measurements* 32(5-6), 787-792.

Pagonis V. and Shannon C. (2000) An improved experimental procedure of separating a composite thermoluminescence glow curve into its components. *Radiation Measurements* **32**(5-6), 805-812.

Rambaud X., Fain J., Miallier D., Pilleyre T., and Sanzelle S. (2000) Annual dose assessment for luminescence and ESR dating: evaluation of nuclides mobility. *Radiation Measurements* 32(5-6), 741-746.

Rhodes E. J. (2000) Observations of thermal transfer OSL signals in glacigenic quartz. Radiation Measurements 32(5-6), 595-602.

Richardson C. A. (2000) Preheat-induced signal enhancement in the infrared stimulated luminescence of young and bleached sediment samples. *Radiation Measurements* 32(5-6), 541-547.

Roberts R. G., Galbraith R. F., Yoshida H., Laslett G. M., and Olley J. M. (2000) Distinguishing dose populations in sediment mixtures: a test of single-grain optical dating procedures using mixtures of laboratory-dosed quartz. *Radiation Measurements* **32**(5-6), 459-465.

Robertson S. and Grun R. (2000) Dose determination on tooth enamel fragments from two human fossils. *Radiation Measurements* **32**(5-6), 773-779.

Schilles T. and Habermann J. (2000) Radioluminescence dating: the IR emission of fefdspar. Radiation Measurements 32(5-6), 679-683.

Sholom S. V., Haskell E. H., Hayes R. B., Chumak V. V., and Kenner G. H. (2000) EPR-dosimetry with carious teeth. *Radiation Measurements* 32(5-6), 799-803.

Short M. A. and Huntley D. J. (2000) Crystal anisotropy effects in optically stimulated luminescence of a K-feldspar. *Radiation Measurements* **32**(5-6), 865-871.

Singarayer J. S., Bailey R. M., and Rhodes E. J. (2000) Potential of the slow component of quartz OSL for age determination of sedimentary samples. *Radiation Measurements* **32**(5-6), 873-880.

Spooner N. A. (2000) A photon-counting imaging system (PCIS) for luminescence applications. *Radiation Measurements* **32**(5-6), 513-521.

Spooner N. A. and Allsop A. (2000) The spatial variation of dose-rate from Sr-90/Y-90 beta sources for use in luminescence dating. *Radiation Measurements* 32(1), 49-56.

Spooner N. A. and Questiaux D. G. (2000) Kinetics of red, blue and UV thermoluminescence and opticallystimulated luminescence from quartz. *Radiation Measurements* **32**(5-6), 659-666.

Stokes S., Colls A. E. L., Fattahi M., and Rich J. (2000) Investigations of the performance of quartz single aliquot D-E determination procedures. *Radiation Measurements* **32**(5-6), 585-594.

Tani A., Yamanaka C., Ikeya M., Ohtaka O., Takada M., and Katsura T. (2000) Optically stimulated luminescence (OSL) study of synthetic stishovite. *Radiation Measurements* **32**(5-6), 473-477.

Toyoda S., Rink W. J., Schwarcz H. P., and Rees-Jones J. (2000) Crushing effects on TL and OSL on quartz: relevance to fault dating. *Radiation Measurements* **32**(5-6), 667-672.

Trautmann T., Krbetschek M. R., Dietrich A., and Stolz W. (2000) The basic principle of radioluminescence dating and a localized transition model. *Radiation Measurements* **32**(5-6), 487-492.

Trautmann T., Krbetschek M. R., and Stolz W. (2000) A systematic study of the radioluminescence properties of single feldspar grains. *Radiation Measurements* **32**(5-6), 685-690.

Truscott A. J., Duller G. A. T., Bøtter-Jensen L., Murray A. S., and Wintle A. G. (2000) Reproducibility of optically stimulated luminescence measurements from single grains of Al2O3 : C and annealed quartz. *Radiation Measurements* **32**(5-6), 447-451.

van Es H. J., den Hartog H. W., de Meijer R. J., Venema L. B., Donoghue J. F., and Rozendaal A. (2000) Assessment of the suitability of zircons for thermoluminescence dating. *Radiation Measurements* 32(5-6), 819-823.

Vandenberghe D., De Corte F., and Van den haute P. (2000) Observations on the thermoluminescence signal induced by alpha radiation. *Radiation Measurements* **32**(5-6), 813-817.

Vartanian E., Guibert P., Rogue C., Bechtel F., and Schvoerer M. (2000) Changes in OSL properties of quartz by preheating: an interpretation. *Radiation Measurements* 32(5-6), 647-652.

Visocekas R. (2000) Monitoring anomalous fading of TL of feldspars by using far-red emission as a gauge. *Radiation Measurements* **32**(5-6), 499-504.

Wallinga J., Murray A., and Duller G. (2000) Underestimation of equivalent dose in single-aliquot optical dating of feldspars caused by preheating. *Radiation Measurements* 32(5-6), 691-695.

Wallinga J., Murray A., and Wintle A. (2000) The single-aliquot regenerative-dose (SAR) protocol applied to coarse-grain feldspar. *Radiation Measurements* **32**(5-6), 529-533.

Wieser A., Mehta K., Amira S., Aragno D., Bercea S., Brik A., Bugai A., Callens F., Chumak V., Ciesielski B., Debuyst R., Dubovsky S., Duliu O. G., Fattibene P., Haskell E. H., Hayes R. B., Ignatiev E. A., Ivannikov A., Kirillov V., Kleschenko E., Nakamura N., Nather M., Nowak J., Onori S., Pass B., Pivovarov S., Romanyukha A., Scherbina O., Shames A. I., Sholom S., Skvortsov V., Stepanenko V., Tikounov D. D., and Toyoda S. (2000) The second international intercomparison on EPR tooth dosimetry. *Radiation Measurements* **32**(5-6), 549-557.

Wintle A. G. and Murray A. S. (2000) Quartz OSL: Effects of thermal treatment and their relevance to laboratory dating procedures. *Radiation Measurements* **32**(5-6), 387-400.

Yoshida H., Roberts R. G., Olley J. M., Laslett G. M., and Galbraith R. F. (2000) Extending the age range of optical dating using single 'supergrains' of quartz. *Radiation Measurements* 32(5-6), 439-446.

Zimprich P. and Hille P. (2000) Investigations of the green- and infrared-stimulated luminescence using the recuperation effect. *Radiation Measurements* **32**(5-6), 535-540.

Zimprich P. and Hille P. (2000) Recuperation of infrared stimulated luminescence of feldspars. Radiation Measurements 32(5-6), 697-701.

Announcements

First announcement

The Fifth World Archaeological Congress

will be held at the Catholic University of America Washington, D.C. USA June 21 - 26, 2003 in partnership with the Smithsonian Institution's National Museum of Natural History & National Museum of the American Indian

** CALL FOR ** SYMPOSIUM PROPOSALS!!

Send to: Joan M. Gero, Academic Secretary Department of Anthropology American University Washington, D.C. 20016 FAX: 202.885.1837 email wac5@american.edu

THEMES INCLUDE (but are not limited to):

Community Archaeologies Consequences of Environmental Change Archaeology for Social Justice Indigenous Arrivals and First Peoples Keeping the Trust: Conservation & Data Management Public Archaeology Archaeology of Recent Time The Politics of Representation: Iconography and Design Social Identities in the Past: Gender, Age, Ethnicity, Class Material Matters: Effective Technologies, Then and Now Archaeology in Practice Archaeology and Cultural Diversity Maritime Archaeology

Congress Patron: Harriet Mayor Fulbright

Congress President: Richard West, Director, National Museum of the American Indian *** http://www.american.edu/wac5 ***

Third International Symposium On Nuclear And related Techniques NURT' 2001

October 22-26, 2001

Havana, Cuba

NURT' 2001 will join several workshops, each devoted to a specific field of research :

- Trace and microanalysis techniques
- Applied radiation physics
- Nuclear techniques for the study and conservation of the cultural heritage
- Radiation technologies
- Applied electronics
- Plant mutation breeding and genetic techniques

http://www.ceaden.cu/nurt_2001/